

***Precalculus A Right Triangle Approach 5th edition  
Ratti Solutions Manual***

SKU: 9780137519415-SOLUTIONS

# INSTRUCTOR'S SOLUTIONS MANUAL

BEVERLY FUSFIELD

## PRECALCULUS A RIGHT TRIANGLE APPROACH FIFTH EDITION

J. S. Ratti

*University of South Florida*

Marcus McWaters

*University of South Florida*

Leslaw A. Skrzypek

*University of South Florida*


Jessica Bernards

*Portland Community College*

Wendy Fresh

*Portland Community College*





The author and publisher of this book have used their best efforts in preparing this book. These efforts include the development, research, and testing of the theories and programs to determine their effectiveness. The author and publisher make no warranty of any kind, expressed or implied, with regard to these programs or the documentation contained in this book. The author and publisher shall not be liable in any event for incidental or consequential damages in connection with, or arising out of, the furnishing, performance, or use of these programs.

Reproduced by Pearson from electronic files supplied by the author.

Copyright © 2023, 2019, 2015 by Pearson Education, Inc. 221 River Street, Hoboken, NJ 07030. All rights reserved.

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written permission of the publisher. Printed in the United States of America.



ISBN-13: 978-0-13-751980-4  
ISBN-10: 0-13-751980-X

# CONTENTS

|                  |                                                                   |     |
|------------------|-------------------------------------------------------------------|-----|
| <b>Chapter P</b> | <b>Basic Concepts of Algebra</b>                                  |     |
| P.1              | The Real Numbers and Their Properties .....                       | 1   |
| P.2              | Integer Exponents and Scientific Notation .....                   | 8   |
| P.3              | Polynomials .....                                                 | 13  |
| P.4              | Factoring Polynomials .....                                       | 20  |
| P.5              | Rational Expressions .....                                        | 28  |
| P.6              | Rational Exponents and Radicals .....                             | 38  |
|                  | Chapter P Review Exercises .....                                  | 48  |
|                  | Chapter P Practice Test .....                                     | 53  |
| <b>Chapter 1</b> | <b>Equations and Inequalities</b>                                 |     |
| 1.1              | Linear Equations in One Variable .....                            | 54  |
| 1.2              | Applications of Linear Equations: Modeling .....                  | 64  |
| 1.3              | Quadratic Equations .....                                         | 76  |
| 1.4              | Complex Numbers: Quadratic Equations with Complex Solutions ..... | 93  |
| 1.5              | Solving Other Types of Equations .....                            | 101 |
| 1.6              | Inequalities .....                                                | 122 |
| 1.7              | Equations and Inequalities Involving Absolute Value .....         | 138 |
|                  | Chapter 1 Key Ideas at a Glance .....                             | 155 |
|                  | Chapter 1 Review Exercises .....                                  | 156 |
|                  | Chapter 1 Practice Test A .....                                   | 167 |
|                  | Chapter 1 Practice Test B .....                                   | 169 |
| <b>Chapter 2</b> | <b>Graphs and Functions</b>                                       |     |
| 2.1              | The Coordinate Plane .....                                        | 171 |
| 2.2              | Graphs of Equations .....                                         | 181 |
| 2.3              | Lines .....                                                       | 194 |
| 2.4              | Functions .....                                                   | 209 |
| 2.5              | Properties of Functions .....                                     | 219 |
| 2.6              | A Library of Functions .....                                      | 229 |
| 2.7              | Transformations of Functions .....                                | 240 |
| 2.8              | Combining Functions; Composite Functions .....                    | 257 |
| 2.9              | Inverse Functions .....                                           | 275 |
|                  | Chapter 2 Key Ideas at a Glance .....                             | 286 |
|                  | Chapter 2 Review Exercises .....                                  | 288 |
|                  | Chapter 2 Practice Test A .....                                   | 298 |
|                  | Chapter 2 Practice Test B .....                                   | 300 |
|                  | Cumulative Review Exercises (Chapters P–2) .....                  | 300 |
| <b>Chapter 3</b> | <b>Polynomial and Rational Functions</b>                          |     |
| 3.1              | Quadratic Functions .....                                         | 305 |
| 3.2              | Polynomial Functions .....                                        | 326 |
| 3.3              | Dividing Polynomials .....                                        | 342 |
| 3.4              | The Real Zeros of a Polynomial Function .....                     | 353 |
| 3.5              | The Complex Zeros of a Polynomial Function .....                  | 376 |
| 3.6              | Rational Functions .....                                          | 386 |
| 3.7              | Variation .....                                                   | 406 |
|                  | Chapter 3 Key Ideas at a Glance .....                             | 411 |
|                  | Chapter 3 Review Exercises .....                                  | 414 |
|                  | Chapter 3 Practice Test A .....                                   | 431 |
|                  | Chapter 3 Practice Test B .....                                   | 432 |

|                                                                 |     |
|-----------------------------------------------------------------|-----|
| Cumulative Review Exercises (Chapters P–3).....                 | 433 |
| <b>Chapter 4 Exponential and Logarithmic Functions</b>          |     |
| 4.1 Exponential Functions.....                                  | 436 |
| 4.2 Logarithmic Functions .....                                 | 449 |
| 4.3 Rules of Logarithms .....                                   | 463 |
| 4.4 Exponential and Logarithmic Equations and Inequalities..... | 475 |
| 4.5 Logarithmic Scales; Modeling .....                          | 490 |
| Chapter 4 Key Ideas at a Glance .....                           | 501 |
| Chapter 4 Review Exercises.....                                 | 501 |
| Chapter 4 Practice Test A.....                                  | 510 |
| Chapter 4 Practice Test B.....                                  | 511 |
| Cumulative Review Exercises (Chapters P–4).....                 | 512 |
| <b>Chapter 5 Trigonometric Functions</b>                        |     |
| 5.1 Angles and Their Measure .....                              | 516 |
| 5.2 Right-Triangle Trigonometry .....                           | 523 |
| 5.3 Trigonometric Functions of Any Angle; The Unit Circle ..... | 535 |
| 5.4 Graphs of the Sine and Cosine Functions.....                | 548 |
| 5.5 Graphs of the Other Trigonometric Functions .....           | 567 |
| 5.6 Inverse Trigonometric Functions .....                       | 582 |
| Chapter 5 Key Ideas at a Glance .....                           | 590 |
| Chapter 5 Review Exercises.....                                 | 592 |
| Chapter 5 Practice Test A.....                                  | 596 |
| Chapter 5 Practice Test B.....                                  | 597 |
| Cumulative Review Exercises (Chapters P–5).....                 | 598 |
| <b>Chapter 6 Trigonometric Identities and Equations</b>         |     |
| 6.1 Verifying Identities .....                                  | 601 |
| 6.2 Sum and Difference Formulas.....                            | 615 |
| 6.3 Double-Angle and Half-Angle Formulas .....                  | 629 |
| 6.4 Product-to-Sum and Sum-to-Product Formulas .....            | 645 |
| 6.5 Trigonometric Equations I.....                              | 654 |
| 6.6 Trigonometric Equations II .....                            | 664 |
| Chapter 6 Key Ideas at a Glance .....                           | 684 |
| Chapter 6 Review Exercises.....                                 | 685 |
| Chapter 6 Practice Test A.....                                  | 691 |
| Chapter 6 Practice Test B.....                                  | 693 |
| Cumulative Review Exercises (Chapters P–6).....                 | 694 |
| <b>Chapter 7 Applications of Trigonometric Functions</b>        |     |
| 7.1 The Law of Sines.....                                       | 698 |
| 7.2 The Law of Cosines.....                                     | 713 |
| 7.3 Areas of Polygons Using Trigonometry.....                   | 726 |
| 7.4 Vectors .....                                               | 741 |
| 7.5 The Dot Product .....                                       | 751 |
| 7.6 Polar Coordinates .....                                     | 761 |
| 7.7 Polar Form of Complex Numbers; DeMoivre’s Theorem .....     | 778 |
| Chapter 7 Key Ideas at a Glance .....                           | 792 |
| Chapter 7 Review Exercises.....                                 | 793 |
| Chapter 7 Practice Test A.....                                  | 802 |
| Chapter 7 Practice Test B.....                                  | 803 |
| Cumulative Review Exercises (Chapters P–7).....                 | 805 |

|                   |                                                      |      |
|-------------------|------------------------------------------------------|------|
| <b>Chapter 8</b>  | <b>Systems of Equations and Inequalities</b>         |      |
| 8.1               | Systems of Linear Equations in Two Variables .....   | 808  |
| 8.2               | Systems of Linear Equations in Three Variables ..... | 828  |
| 8.3               | Partial-Fraction Decomposition .....                 | 851  |
| 8.4               | Systems of Nonlinear Equations .....                 | 875  |
| 8.5               | Systems of Inequalities.....                         | 889  |
| 8.6               | Linear Programming.....                              | 905  |
|                   | Chapter 8 Key Ideas at a Glance .....                | 918  |
|                   | Chapter 8 Review Exercises .....                     | 918  |
|                   | Chapter 8 Practice Test A.....                       | 931  |
|                   | Chapter 8 Practice Test B.....                       | 934  |
|                   | Cumulative Review Exercises (Chapters P–8).....      | 937  |
| <b>Chapter 9</b>  | <b>Matrices and Determinants</b>                     |      |
| 9.1               | Matrices and Systems of Equations.....               | 940  |
| 9.2               | Matrix Algebra .....                                 | 963  |
| 9.3               | The Matrix Inverse .....                             | 983  |
| 9.4               | Determinants and Cramer’s Rule .....                 | 1004 |
|                   | Chapter 9 Key Ideas at a Glance .....                | 1015 |
|                   | Chapter 9 Review Exercises .....                     | 1018 |
|                   | Chapter 9 Practice Test A.....                       | 1029 |
|                   | Chapter 9 Practice Test B.....                       | 1030 |
|                   | Cumulative Review Exercises (Chapters P–9).....      | 1032 |
| <b>Chapter 10</b> | <b>The Conic Sections</b>                            |      |
| 10.2              | The Parabola.....                                    | 1034 |
| 10.3              | The Ellipse.....                                     | 1050 |
| 10.4              | The Hyperbola.....                                   | 1069 |
|                   | Chapter 10 Key Ideas at a Glance .....               | 1092 |
|                   | Chapter 10 Review Exercises.....                     | 1093 |
|                   | Chapter 10 Practice Test A.....                      | 1102 |
|                   | Chapter 10 Practice Test B.....                      | 1104 |
|                   | Cumulative Review Exercises (Chapters P–10).....     | 1105 |
| <b>Chapter 11</b> | <b>Further Topics in Algebra</b>                     |      |
| 11.1              | Sequences and Series .....                           | 1108 |
| 11.2              | Arithmetic Sequences; Partial Sums .....             | 1120 |
| 11.3              | Geometric Sequences and Series.....                  | 1127 |
| 11.4              | Mathematical Induction.....                          | 1138 |
| 11.5              | The Binomial Theorem .....                           | 1149 |
| 11.6              | Counting Principles .....                            | 1157 |
| 11.7              | Probability .....                                    | 1164 |
|                   | Chapter 11 Key Ideas at a Glance .....               | 1168 |
|                   | Chapter 11 Review Exercises .....                    | 1169 |
|                   | Chapter 11 Practice Test A.....                      | 1173 |
|                   | Chapter 11 Practice Test B.....                      | 1173 |
|                   | Cumulative Review Exercises (Chapters P–11).....     | 1174 |



# Chapter P Basic Concepts of Algebra

## P.1 The Real Numbers and Their Properties

### Practice Problems


- Let  $x = 2.132132132\dots$   
Then,  $1000x = 2132.132132\dots$   
 $1000x = 2132.132132\dots$   
 $x = 2.132132\dots$   

---

 $999x = 2130$   
 $x = \frac{2130}{999} = \frac{710}{333}$
- Natural numbers: 2, 7
  - Whole numbers: 0, 2, 7
  - Integers:  $-6, -\frac{21}{7} = -3, 0, 2, 7$
  - Rational numbers:  
 $-6, -\frac{21}{7} (= -3), -\frac{1}{2}, 0, \frac{4}{3}, 2, 7$
  - Irrational numbers:  $\sqrt{3}, \sqrt{17}$
  - Real numbers: the set B
- $2^3 = 2 \cdot 2 \cdot 2 = 8$
  - $(3a)^2 = (3a)(3a) = 9a^2$
  - $\left(\frac{1}{2}\right)^4 = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{16}$
- $-2 < 0$  true because  $-2$  is to the left of 0 on the number line.
  - $5 \leq 7$  true because 5 is to the left of 7 on the number line.
  - $-4 > -1$  false because  $-4$  is to the left of  $-1$  on the number line.
- $A = \{-3, -1, 0, 1, 3\}, B = \{-4, -2, 0, 2, 4\}$   
 $A \cap B = \{0\}$   
 $A \cup B = \{-4, -3, -2, -1, 0, 1, 2, 3, 4\}$
- $I_1 \cup I_2 = (-\infty, \infty)$
  - $I_1 \cap I_2 = [-2, 5)$
- $|-10| = 10$

$$\text{b. } |3 - 4| = |-1| = 1$$

$$\text{c. } |2(-3) + 7| = |-6 + 7| = |1| = 1$$



$$d(-7, 2) = |-7 - 2| = |-9| = 9$$

$$\text{9. a. } (-3) \cdot 5 + 20 = -15 + 20 = 5$$

$$\text{b. } 5 - 12 \div 6 \cdot 2 = 5 - 2 \cdot 2 = 5 - 4 = 1$$

$$\text{c. } \frac{9-1}{4} - 5 \cdot 7 = \frac{8}{4} - 5 \cdot 7 = 2 - 35 = -33$$

$$\text{d. } -3 + (x-4)^2 \text{ for } x = 6$$

$$-3 + (6-4)^2 = -3 + 2^2 = -3 + 4 = 1$$

$$\text{10. a. } \frac{7}{4} + \frac{3}{8} = \frac{7 \cdot 2}{4 \cdot 2} + \frac{3}{8} = \frac{7 \cdot 2 + 3}{8} = \frac{17}{8}$$

$$\text{b. } \frac{8}{3} - \frac{2}{5} = \frac{8 \cdot 5}{3 \cdot 5} - \frac{2 \cdot 3}{5 \cdot 3} = \frac{8 \cdot 5 - 2 \cdot 3}{3 \cdot 5} = \frac{34}{15}$$

$$\text{c. } \frac{9}{14} \cdot \frac{7}{3} = \frac{9 \cdot 7}{14 \cdot 3} = \frac{3 \cdot \cancel{7} \cdot \cancel{7}}{2 \cdot \cancel{7} \cdot \cancel{3}} = \frac{3}{2}$$

$$\text{d. } \frac{\frac{5}{8}}{\frac{15}{16}} = \frac{5}{8} \cdot \frac{16}{15} = \frac{5 \cdot 16}{8 \cdot 15} = \frac{\cancel{5} \cdot 2 \cdot \cancel{8}}{\cancel{8} \cdot 3 \cdot \cancel{3}} = \frac{2}{3}$$

$$\text{11. a. } (x-2) \div 3 + x = (3-2) \div 3 + 3$$

$$= 1 \div 3 + 3 = \frac{1}{3} + 3 = \frac{10}{3}$$

$$\text{b. } 7 - \frac{x}{|y|} = 7 - \frac{-1}{|3|} = 7 + \frac{1}{3} = \frac{22}{3}$$

$$\text{12. } ^\circ\text{C} = \frac{39}{3} + 4 = 13 + 4 = 17^\circ\text{C}$$

### Concepts and Vocabulary

- Whole numbers are formed by adding the number zero to the set of natural numbers.
- The number  $-3$  is an integer, but it is also a rational number and a real number.
- If  $a < b$ , then  $a$  is to the left of  $b$  on the number line.

4. If a real number is not a rational number, it is an irrational number.
5. True
6. False.  $\frac{-5}{2} = -2\frac{1}{2}$
7. True
8. False. An example is  $(2 + \sqrt{2}) + (2 - \sqrt{2}) = 4$ , which is rational.

**Building Skills**

9.  $0.\bar{3}$ , repeating
10.  $0.\bar{6}$ , repeating
11.  $-0.8$ , terminating
12.  $-0.25$ , terminating
13.  $0.\bar{27}$ , repeating
14.  $0.\bar{3}$ , repeating
15.  $3.\bar{16}$ , repeating
16.  $2.\bar{73}$ , repeating
17.  $3.75 = \frac{375}{100} = \frac{15}{4}$
18.  $-2.35 = -\frac{235}{100} = -\frac{47}{20}$
19.  $10x = 53.\bar{3}$   
 $\frac{x = 5.\bar{3}}{9x = 48}$   
 $x = \frac{48}{9} = \frac{16}{3}$   
 $-5.\bar{3} = -\frac{16}{3}$
20.  $10x = 96.\bar{6}$   
 $\frac{x = 9.\bar{6}}{9x = 87}$   
 $x = \frac{87}{9} = \frac{29}{3}$
21.  $100x = 213.\bar{13}$   
 $\frac{x = 2.\bar{13}}{99x = 211}$   
 $x = \frac{211}{99}$
22.  $100x = 323.\bar{23}$   
 $\frac{x = 3.\bar{23}}{99x = 320}$   
 $x = \frac{320}{99}$
23.  $100x = 452.3\bar{23}$   
 $\frac{x = 4.5\bar{23}}{99x = 447.8}$   
 $x = \frac{447.8}{99} = \frac{4478}{990} = \frac{2239}{495}$

$$24. \begin{array}{r} 100x = 142.3\bar{535} \\ x = 1.4\bar{235} \\ \hline 99x = 140.93 \\ x = \frac{140.93}{99} = \frac{14,093}{9900} \end{array}$$

25. Rational                      26. Rational
27. Rational                      28. Rational
29. Rational                      30. Rational
31. Irrational                      32. Irrational

Exercises 33-38 refer to the set


$$A = \left\{ -19, -\frac{12}{3}, -\sqrt{3}, 0, 2, \sqrt{10}, \frac{17}{4}, 11 \right\}$$

33. Natural numbers: 2, 11
34. Whole numbers: 0, 2, 11
35. Integers:  $-19, -\frac{12}{3} = -4, 0, 2, 11$
36. Rational numbers:  $-19, -\frac{12}{3}, 0, 2, \frac{17}{4}, 11$
37. Irrational numbers:  $\sqrt{3}, \sqrt{10}$
38. Real numbers: All numbers in set  $A$  are real numbers.
39.  $10^3$   
 base: 10; exponent: 3  
 $10^3 = 10 \cdot 10 \cdot 10 = 1000$
40.  $5^4$   
 base: 5; exponent: 4  
 $5^4 = 5 \cdot 5 \cdot 5 \cdot 5 = 625$
41.  $\left(\frac{2}{3}\right)^3$   
 base:  $\frac{2}{3}$ ; exponent: 3  
 $\left(\frac{2}{3}\right)^3 = \frac{2}{3} \cdot \frac{2}{3} \cdot \frac{2}{3} = \frac{8}{27}$
42.  $\left(\frac{5}{2}\right)^4$   
 base:  $\frac{5}{2}$ ; exponent: 4  
 $\left(\frac{5}{2}\right)^4 = \frac{5}{2} \cdot \frac{5}{2} \cdot \frac{5}{2} \cdot \frac{5}{2} = \frac{625}{16}$


43.  $(-2)^3$   
base:  $-2$ ; exponent: 3  
 $(-2)^3 = (-2) \cdot (-2) \cdot (-2) = -8$
44.  $\left(-\frac{1}{2}\right)^4$   
base:  $-\frac{1}{2}$ ; exponent: 4  
 $\left(-\frac{1}{2}\right)^4 = \left(-\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{1}{2}\right) = \frac{1}{16}$
45.  $2 \cdot 3^4$   
base: 3; exponent: 4  
 $2 \cdot 3^4 = 2 \cdot 3 \cdot 3 \cdot 3 \cdot 3 = 2 \cdot 81 = 162$
46.  $5 \cdot \left(\frac{1}{3}\right)^3$   
base:  $\frac{1}{3}$ ; exponent: 3  
 $5 \cdot \left(\frac{1}{3}\right)^3 = 5 \cdot \frac{1}{3} \cdot \frac{1}{3} \cdot \frac{1}{3} = 5 \cdot \frac{1}{27} = \frac{5}{27}$
47.  $-2 \cdot 3^4$   
base: 3; exponent: 4  
 $-2 \cdot 3^4 = -2 \cdot 3 \cdot 3 \cdot 3 \cdot 3 = -2 \cdot 81 = -162$
48.  $-3 \cdot (-2)^4$   
base:  $-2$ ; exponent: 4  
 $-3 \cdot (-2)^4 = -3(-2)(-2)(-2)(-2) = -3 \cdot 16 = -48$
49.  $-3 \cdot (-2)^5$   
base:  $-2$ ; exponent: 5  
 $-3 \cdot (-2)^5 = -3(-2)(-2)(-2)(-2)(-2) = -3 \cdot (-32) = 96$
50.  $-5 \cdot (-3)^2$   
base:  $-3$ ; exponent: 2  
 $-5 \cdot (-3)^2 = -5(-3)(-3) = -5 \cdot 9 = -45$
51.  $3 > -2$
52.  $-3 < -2$
53.  $\frac{1}{2} \geq \frac{1}{2}$
54.  $x < x + 1$
55.  $5 \leq 2x$
56.  $x - 1 > 2$
57.  $-x > 0$
58.  $x < 0$
59.  $2x + 7 \leq 14$
60.  $2x + 3 \leq 5$
61.  $4 = \frac{24}{6}$
62.  $-5 < -2$
63.  $-4 < 0$
64.  $-\frac{9}{2} = -4\frac{1}{2}$
65.  $A \cup B = \{-4, -3, -2, 0, 1, 2, 3, 4\}$
66.  $A \cap B = \{0, 2, 4\}$
67.  $A \cap C = \{-4, -2, 0, 2\}$
68.  $B \cup C = \{-4, -3, -2, -1, 0, 1, 2, 3, 4\}$
69.  $(B \cap C) \cup A = \{-3, 0, 2\} \cup A = \{-4, -3, -2, 0, 2, 4\}$
70.  $(A \cup C) \cap B = \{-4, -3, -2, -1, 0, 2, 4\} \cap B = \{-3, 0, 2, 4\}$
71.  $(A \cup B) \cap C = \{-4, -3, -2, 0, 1, 2, 3, 4\} \cap C = \{-4, -3, -2, 0, 2\}$
72.  $(A \cup B) \cup C = \{-4, -3, -2, 0, 1, 2, 3, 4\} \cup C = \{-4, -3, -2, -1, 0, 1, 2, 3, 4\}$
73.  $I_1 \cup I_2 = (-2, 5)$ ;  $I_1 \cap I_2 = [1, 3]$
74.  $I_1 \cup I_2 = [1, 7]$ ;  $I_1 \cap I_2 = (3, 5)$
75.  $I_1 \cup I_2 = (-6, 10)$ ;  $I_1 \cap I_2 = \emptyset$
76.  $I_1 \cup I_2 = (-\infty, \infty)$ ;  $I_1 \cap I_2 = \emptyset$
77.  $I_1 \cup I_2 = (-\infty, \infty)$ ;  $I_1 \cap I_2 = [2, 5)$
78.  $I_1 \cup I_2 = (-2, \infty)$ ;  $I_1 \cap I_2 = (0, \infty)$
79.  $I_1 \cup I_2 = (-\infty, \infty)$ ;  $I_1 \cap I_2 = [-1, 3) \cup [5, 7]$
80.  $I_1 \cup I_2 = (-\infty, 2] \cup (6, \infty)$ ;  $I_1 \cap I_2 = [-3, 0]$
81.  $|-4| = 4$
82.  $-|-17| = -17$
83.  $\left|\frac{5}{-7}\right| = \frac{5}{7}$
84.  $\left|\frac{-3}{5}\right| = \frac{3}{5}$
85.  $|5 - \sqrt{2}| = 5 - \sqrt{2}$
86.  $|\sqrt{2} - 5| = 5 - \sqrt{2}$
87.  $|\sqrt{3} - 2| = 2 - \sqrt{3}$
88.  $|3 - \pi| = \pi - 3$
89.  $\frac{8}{|-8|} = \frac{8}{8} = 1$
90.  $\frac{-8}{|8|} = \frac{-8}{8} = -1$

91.  $|5 - |-7|| = |5 - 7| = |-2| = 2$


92.  $||4| - 7|| = |4 - 7| = |-3| = 3$




$$d(3, 8) = |3 - 8| = |-5| = 5$$




$$d(2, 14) = |2 - 14| = |-12| = 12$$




$$d(-6, 9) = |-6 - 9| = |-15| = 15$$




$$d(-12, 3) = |-12 - 3| = |-15| = 15$$




$$d(-20, -6) = |-20 - (-6)| = |-14| = 14$$




$$d(-14, -1) = |-14 - (-1)| = |-13| = 13$$



$$d\left(\frac{22}{7}, -\frac{4}{7}\right) = \left|\frac{22}{7} - \left(-\frac{4}{7}\right)\right| = \left|\frac{26}{7}\right| = \frac{26}{7}$$



$$d\left(\frac{16}{5}, -\frac{3}{5}\right) = \left|\frac{16}{5} - \left(-\frac{3}{5}\right)\right| = \left|\frac{19}{5}\right| = \frac{19}{5}$$



$$-3 < x \leq 1$$



$$-6 \leq x < -2$$




$$x \geq -3$$



$$x \geq 0$$



$$x \leq 5$$



$$x \leq -1$$



$$-\frac{3}{4} < x < \frac{9}{4}$$



$$-3 < x < -\frac{1}{2}$$

109.  $4(x + 1) = 4x + 4$

110.  $(-3)(2 - x) = -6 + 3x$

111.  $5(x - y + 1) = 5x - 5y + 5$

112.  $2(3x + 5 - y) = 6x + 10 - 2y$

113. Additive inverse:  $-5$ ; reciprocal:  $\frac{1}{5}$

114. Additive inverse:  $\frac{2}{3}$ ; reciprocal:  $-\frac{3}{2}$

115. Additive inverse:  $0$ ; no reciprocal

116. Additive inverse:  $-1.7$ ; reciprocal:  $\frac{10}{17}$

117. Additive inverse

118. Additive inverse

119. Multiplicative identity

120. Multiplicative identity

121. Associative property of multiplication

122. Associative property of multiplication

123. Multiplicative inverse

124. Multiplicative inverse

125. Additive identity

126. Additive identity

127. Associative property of addition

128. Commutative and associative properties of addition

$$129. \frac{3}{5} + \frac{4}{3} = \frac{3 \cdot 3}{5 \cdot 3} + \frac{4 \cdot 5}{3 \cdot 5} = \frac{9 + 20}{15} = \frac{29}{15}$$

$$130. \frac{7}{10} + \frac{3}{4} = \frac{7 \cdot 2}{10 \cdot 2} + \frac{3 \cdot 5}{4 \cdot 5} = \frac{14 + 15}{20} = \frac{29}{20}$$

$$131. \frac{6 \cdot 7}{5 \cdot 7} + \frac{5 \cdot 5}{7 \cdot 5} = \frac{42 + 25}{35} = \frac{67}{35}$$

$$132. \frac{9}{2} + \frac{5}{12} = \frac{9 \cdot 6}{2 \cdot 6} + \frac{5}{12} = \frac{54 + 5}{12} = \frac{59}{12}$$

$$133. \frac{5}{6} + \frac{3}{10} = \frac{5 \cdot 5}{6 \cdot 5} + \frac{3 \cdot 3}{10 \cdot 3} = \frac{25 + 9}{30} = \frac{34}{30} = \frac{17}{15}$$

$$134. \frac{8}{15} + \frac{2}{9} = \frac{8 \cdot 3}{15 \cdot 3} + \frac{2 \cdot 5}{9 \cdot 5} = \frac{24 + 10}{45} = \frac{34}{45}$$

$$135. \frac{5}{8} - \frac{9}{10} = \frac{5 \cdot 5}{8 \cdot 5} - \frac{9 \cdot 4}{10 \cdot 4} = \frac{25 - 36}{40} = -\frac{11}{40}$$

$$136. \frac{7}{8} - \frac{1}{5} = \frac{7 \cdot 5}{8 \cdot 5} - \frac{1 \cdot 8}{5 \cdot 8} = \frac{35 - 8}{40} = \frac{27}{40}$$

$$137. \frac{5}{9} - \frac{7}{11} = \frac{5 \cdot 11}{9 \cdot 11} - \frac{7 \cdot 9}{11 \cdot 9} = \frac{55 - 63}{99} = -\frac{8}{99}$$

$$138. \frac{5}{8} - \frac{7}{12} = \frac{5 \cdot 3}{8 \cdot 3} - \frac{7 \cdot 2}{12 \cdot 2} = \frac{15 - 14}{24} = \frac{1}{24}$$

$$139. \frac{2}{5} - \frac{1}{2} = \frac{2 \cdot 2}{5 \cdot 2} - \frac{1 \cdot 5}{2 \cdot 5} = \frac{4 - 5}{10} = -\frac{1}{10}$$

$$140. \frac{1}{4} - \frac{1}{6} = \frac{1 \cdot 3}{4 \cdot 3} - \frac{1 \cdot 2}{6 \cdot 2} = \frac{3 - 2}{12} = \frac{1}{12}$$

$$141. \frac{3}{4} \cdot \frac{8}{27} = \frac{3 \cdot 8}{4 \cdot 27} = \frac{\cancel{3} \cdot 2 \cdot \cancel{4}}{\cancel{4} \cdot \cancel{3} \cdot 9} = \frac{2}{9}$$

$$142. \frac{9}{7} \cdot \frac{14}{27} = \frac{9 \cdot 14}{7 \cdot 27} = \frac{\cancel{9} \cdot 2 \cdot \cancel{7}}{\cancel{7} \cdot 3 \cdot \cancel{9}} = \frac{2}{3}$$

$$143. \frac{\frac{8}{5}}{\frac{16}{15}} = \frac{8}{5} \cdot \frac{15}{16} = \frac{8 \cdot 15}{5 \cdot 16} = \frac{\cancel{8} \cdot 3 \cdot \cancel{5}}{\cancel{5} \cdot 2 \cdot \cancel{8}} = \frac{3}{2}$$

$$144. \frac{\frac{5}{6}}{\frac{15}{6}} = \frac{5}{6} \cdot \frac{6}{15} = \frac{5 \cdot 6}{6 \cdot 15} = \frac{\cancel{6} \cdot \cancel{6}}{\cancel{6} \cdot 3 \cdot \cancel{5}} = \frac{1}{3}$$

$$145. \frac{\frac{7}{8}}{\frac{21}{16}} = \frac{7}{8} \cdot \frac{16}{21} = \frac{7 \cdot 16}{8 \cdot 21} = \frac{\cancel{7} \cdot 2 \cdot \cancel{8}}{\cancel{8} \cdot 3 \cdot \cancel{7}} = \frac{2}{3}$$

$$146. \frac{\frac{3}{10}}{\frac{7}{15}} = \frac{3}{10} \cdot \frac{15}{7} = \frac{3 \cdot 15}{10 \cdot 7} = \frac{3 \cdot 3 \cdot \cancel{5}}{2 \cdot \cancel{5} \cdot 7} = \frac{9}{14}$$

$$147. 5 \cdot \frac{3}{10} - \frac{1}{2} = \frac{\cancel{5} \cdot 3}{1 \cdot 2 \cdot \cancel{5}} - \frac{1}{2} = \frac{3}{2} - \frac{1}{2} = \frac{2}{2} = 1$$

$$148. 2 \cdot \frac{7}{2} - \frac{3}{2} = \frac{14}{2} - \frac{3}{2} = \frac{11}{2}$$

$$149. 3 \cdot \frac{2}{15} - \frac{1}{3} = \frac{3 \cdot 2}{1 \cdot 15} - \frac{1}{3} = \frac{\cancel{3} \cdot 2}{1 \cdot \cancel{3} \cdot 5} - \frac{1}{3} = \frac{2}{5} - \frac{1}{3} = \frac{2 \cdot 3}{5 \cdot 3} - \frac{1 \cdot 5}{3 \cdot 5} = \frac{6}{15} - \frac{5}{15} = \frac{1}{15}$$

$$150. 2 \cdot \frac{5}{3} - \frac{3}{2} = \frac{10}{3} - \frac{3}{2} = \frac{20}{6} - \frac{9}{6} = \frac{11}{6}$$

$$151. 2(x + y) - 3y = 2(3 + (-5)) - 3(-5) = 2(-2) - (-15) = -4 + 15 = 11$$

$$152. -2(x + y) + 5y = -2(3 + (-5)) + 5(-5) = -2(-2) + (-25) = 4 + (-25) = -21$$

$$153. 3|x| - 2|y| = 3|3| - 2|-5| = 3(3) - 2(5) = 9 - 10 = -1$$

$$154. 7|x - y| = 7|3 - (-5)| = 7|8| = 7(8) = 56$$

$$155. \frac{x - 3y}{2} + xy = \frac{3 - 3(-5)}{2} + 3(-5) = \frac{3 - (-15)}{2} + (-15) = \frac{18}{2} + (-15) = 9 + (-15) = -6$$

$$156. \frac{y+3}{x} - xy = \frac{-5+3}{3} - 3(-5) \\ = \frac{-2}{3} - (-15) = \frac{43}{3}$$

$$157. \frac{2(1-2x)}{y} - (-x)y = \frac{2(1-2(3))}{-5} - (-3)(-5) \\ = \frac{2(-5)}{-5} - 15 = 2 - 15 = -13$$

$$158. \frac{3(2-x)}{y} - (1-xy) = \frac{3(2-3)}{-5} - (1-3(-5)) \\ = \frac{3(-1)}{-5} - (1-(-15)) \\ = \frac{3}{5} - 16 = -\frac{77}{5}$$

$$159. \frac{\frac{14}{x} + \frac{1}{2}}{\frac{-y}{4}} = \frac{\frac{14}{3} + \frac{1}{2}}{\frac{-(-5)}{4}} = \frac{31}{6} \cdot \frac{4}{5} = \frac{62}{15}$$

$$160. \frac{\frac{4}{-y} + \frac{8}{x}}{\frac{y}{2}} = \frac{\frac{4}{-(-5)} + \frac{8}{3}}{\frac{-5}{2}} = \left(\frac{4}{5} + \frac{8}{3}\right) \cdot \left(-\frac{2}{5}\right) \\ = \frac{52}{15} \cdot \left(-\frac{2}{5}\right) = -\frac{104}{75}$$

$$161. \frac{x}{y} + \frac{x}{3} = \frac{3x+xy}{3y}$$

$$162. (x+2)(x+3) = x^2 + 3x + 2x + 6 = x^2 + 5x + 6$$

$$163. 5(x+3) = 5x + 5(3) = 5x + 15$$

$$164. (25x)(4x) = 100x^2$$

$$165. x - (3y + 2) = x - 3y - 2$$

$$166. 2x - (4y - 5) = 2x - 4y + 5$$

$$167. \frac{x+y}{x} = \frac{x}{x} + \frac{y}{x} = 1 + \frac{y}{x}$$

$$168. \frac{x+y}{x+z} = \frac{x}{x+z} + \frac{y}{x+z}$$

$$169. (x+1)(y+1) = xy + x + y + 1$$

$$170. \frac{\frac{x}{1}}{\frac{1}{y}} = \frac{1}{x} \cdot \frac{y}{1} = \frac{y}{x}$$

$$171. d(P, M) = \left| \frac{a+b}{2} - a \right| = \left| \frac{a+b-2a}{2} \right| = \left| \frac{-a+b}{2} \right| \\ d(Q, M) = \left| b - \frac{a+b}{2} \right| = \left| \frac{2b-a-b}{2} \right| = \left| \frac{b-a}{2} \right|$$

Since

$$d(P, M) = \left| \frac{-a+b}{2} \right| = \left| \frac{b-a}{2} \right| = d(Q, M), M \text{ is} \\ \text{the midpoint of the line segment } PQ.$$

172. Answers may vary. Using the hint, we have

$$-\frac{4}{13} = -\frac{40}{130} \text{ and } \frac{7}{13} = \frac{70}{130}. \text{ Therefore,} \\ -\frac{40}{130} < -\frac{31}{130} < -\frac{30}{130} < \dots < 0 < \dots \\ < \frac{1}{130} < \frac{10}{130} < \dots < \frac{69}{130} < \frac{70}{130}.$$

### Applying the Concepts

173. a. people who own either MP3 players or people who own DVD players.

b. people who own both MP3 players and DVD players.

174. a.  $A = \{\text{Ford Bronco, Honda Accord}\}$

b.  $B = \{\text{Ford Bronco}\}$

c.  $C = \{\text{Ford Bronco, Honda Accord, Audi A4}\}$

d.  $A \cap B = \{\text{Ford Bronco}\}$

e.  $B \cap C = \{\text{Ford Bronco}\}$

f.  $A \cup B = \{\text{Ford Bronco, Honda Accord}\}$

g.  $A \cup C = \{\text{Ford Bronco, Honda Accord, Audi A4}\}$

175.  $119.5 \leq x \leq 134.5$



176.  $49 \leq x \leq 95$



177. a.  $|124 - 120| = 4$

b.  $|137 - 120| = 17$

c.  $|114 - 120| = |-6| = 6$

178. a.  $|72 - 56| = 16$  MB per hour

b.  $|700 - 380| = 320$  MB per hour

179.  $^{\circ}\text{C} = \frac{42}{3} + 4 = 14 + 4 = 18^{\circ}\text{C}$

180.  $22 = \frac{c}{3} + 4 \Rightarrow 18 = \frac{c}{3} \Rightarrow c = 54$

If the Celsius temperature is  $22^{\circ}$ , then 54 chirps will be counted.

181. Let  $x$  = the number of calories from broccoli.  
Then we have  
 $522.5 - 55x = 0 \Rightarrow 522.5 = 55x \Rightarrow 9.5 = x$   
The number of grams of broccoli is  
 $9.5 \times 100 = 950$  grams.

182. Let  $x$  = the number of orders of french fries.  
The number of calories lost from broccoli is  
 $6 \times 55 = 330$ . Then we have  
 $165x - 330 = 0 \Rightarrow 165x = 330 \Rightarrow x = 2$   
So, Carmen will have to eat 2 orders of french fries.

### Beyond the Basics

183. True                      184. True

185. a. False. For example,  $0 \cdot \sqrt{2} = 0$ .

b. The products of a nonzero rational number and an irrational number is an irrational number.

186. False. For example,  $\sqrt{2} \cdot \sqrt{2} = 2$ .

187. False. For example,  $(2 + \sqrt{3}) - (6 + \sqrt{3}) = -4$ .

188. False. For example,  $\frac{\sqrt{12}}{\sqrt{3}} = \sqrt{4} = 2$ .

189. True                      190. True

For exercises 191–192, use the following definition:  
An integer  $P$  is *even* if  $p = 2n$  for some integer  $n$ . An integer  $q$  is *odd* if  $q = 2k + 1$  for some integer  $k$ .

191. a. If  $a$  is odd, then  $a = 2m + 1$  for some integer  $m$ .

$$\begin{aligned} a^2 &= (2m+1)^2 = (2m+1)(2m+1) \\ &= 4m^2 + 4m + 1 = 4m(m+1) + 1 \\ &= 2(2m(m+1)) + 1, \end{aligned}$$

which is of the format  $2k + 1$ . Therefore,  $a^2$  is an odd integer.

b. Note that this statement is the contrapositive of the statement in part (a). That is, if the statement is “if  $p$ , then  $q$ ”, the contrapositive is “if not  $q$ , then not  $p$ .”

The contrapositive of “If an integer  $a$  is odd, then  $a^2$  is also an odd integer” is “If  $a^2$  is not an odd integer (i.e., an even integer), then  $a$  is also not an odd integer (i.e., an even integer.” Logically, these two statements are true. That means that the original statement and its contrapositive are either both true or both false. We already know from part (a) that the original statement is true. Therefore, the contrapositive is also true. Thus, if  $b^2$  is even, then  $b$  is also an even integer.

192.  $2q^2$  is even because it is of the form  $2n$ , so it follows that  $p^2$  is even. Then, from exercise 191(b),  $p$  is also even. Therefore,  $p = 2n$ , for some integer  $n$ . Substituting, we have  
 $2q^2 = p^2 = (2n)^2 \Rightarrow 2q^2 = 2 \cdot 2n^2 \Rightarrow$   
 $q^2 = 2n^2$ , and, thus,  $q$  is even using exercise 191 (b).

### Getting Ready for the Next Section

GR1. a.  $a^2 \cdot a^3 = a^5$

b.  $a^4 \cdot a^7 = a^{11}$

c.  $a^m \cdot a^n = a^{m+n}$

GR2. a.  $\frac{b^3}{b} = b^2$

b.  $\frac{b^7}{b^3} = b^4$

c.  $\frac{b^m}{b^n} = b^{m-n}$

GR3. a.  $(a^2)^3 = a^6$

b.  $(a^4)^2 = a^8$

c.  $(a^m)^n = a^{m \cdot n} = a^{mn}$

GR4. a.  $(ab)^2 = a^2 b^2$

b.  $(ab)^4 = a^4 b^4$

c.  $(ab)^n = a^n b^n$

$$\text{GR5. } \frac{7}{3} + \frac{3}{2} = \frac{7 \cdot 2}{3 \cdot 2} + \frac{3 \cdot 3}{2 \cdot 3} = \frac{14 + 9}{6} = \frac{23}{6}$$

$$\text{GR6. } \frac{3}{6} - \frac{2}{3} = \frac{3}{6} - \frac{2 \cdot 2}{3 \cdot 2} = \frac{3 - 4}{6} = -\frac{1}{6}$$

$$\text{GR7. } \frac{3}{2} \cdot \frac{12}{7} = \frac{3 \cdot 12}{2 \cdot 7} = \frac{3 \cdot \cancel{2} \cdot 6}{\cancel{2} \cdot 7} = \frac{18}{7}$$

$$\text{GR8. } \frac{3}{5} \div \frac{12}{15} = \frac{3}{5} \cdot \frac{15}{12} = \frac{3 \cdot 15}{5 \cdot 12} = \frac{\cancel{3} \cdot 3 \cdot \cancel{5}}{\cancel{5} \cdot \cancel{3} \cdot 4} = \frac{3}{4}$$

## P.2 Integer Exponents and Scientific Notation

### Practice Exercises

$$1. \text{ a. } 2^{-1} = \frac{1}{2}$$

$$\text{b. } \left(\frac{4}{5}\right)^0 = 1$$

$$\text{c. } \left(\frac{3}{2}\right)^{-2} = \frac{1}{(3/2)^2} = \frac{1}{9/4} = \frac{4}{9}$$

$$2. \text{ a. } x^2 \cdot 3x^7 = 3x^9$$

$$\text{b. } (2^2 x^3)(4x^{-3}) = 4x^3 \cdot 4x^{-3} = 16x^{3+(-3)} = 16x^0 = 16 \cdot 1 = 16$$

$$3. \text{ a. } \frac{3^4}{3^0} = 3^{4-0} = 3^4 = 81$$

$$\text{b. } \frac{5}{5^{-2}} = 5^{1-(-2)} = 5^3 = 125$$

$$\text{c. } \frac{2x^3}{3x^{-4}} = \frac{2x^{3-(-4)}}{3} = \frac{2x^7}{3}$$

$$4. \text{ a. } (7^{-5})^0 = 7^{(-5)(0)} = 7^0 = 1$$

$$\text{b. } (7^0)^{-5} = 7^{(0)(-5)} = 7^0 = 1$$

$$\text{c. } (x^{-1})^8 = x^{(-1)(8)} = x^{-8} = \frac{1}{x^8}$$

$$\text{d. } (x^{-2})^{-5} = x^{(-2)(-5)} = x^{10}$$

$$5. \text{ a. } \left(\frac{1}{2}x\right)^{-1} = \left(\frac{1}{2}\right)^{-1} (x)^{-1} = 2\left(\frac{1}{x}\right) = \frac{2}{x}$$

$$\begin{aligned} \text{b. } (5x^{-1})^2 &= 5^2 (x^{-1})^2 = 5^2 x^{(-1)(2)} \\ &= 25x^{-2} = \frac{25}{x^2} \end{aligned}$$

$$\text{c. } (xy^2)^3 = x^3 y^{2(3)} = x^3 y^6$$

$$\text{d. } (x^{-2}y)^{-3} = x^{(-2)(-3)} y^{-3} = \frac{x^6}{y^3}$$

$$6. \text{ a. } \left(\frac{1}{3}\right)^2 = \frac{1^2}{3^2} = \frac{1}{9}$$

$$\text{b. } \left(\frac{10}{7}\right)^{-2} = \left(\frac{7}{10}\right)^2 = \frac{7^2}{10^2} = \frac{49}{100}$$

$$7. \text{ a. } (2x^4)^{-2} = \left(\frac{1}{2x^4}\right)^2 = \frac{1}{4x^8}$$

$$\text{b. } \frac{x^2(-y)^3}{(xy^2)^3} = \frac{-x^2y^3}{x^3y^6} = -\frac{1}{xy^3}$$

$$8. \quad 732,000 = 7.32 \times 10^5$$

$$\begin{aligned} 9. \quad \frac{2.1 \times 10^{10}}{3.5 \times 10^8} &= \frac{2.1}{3.5} \times \frac{10^{10}}{10^8} = \frac{2.10}{3.5} \times 10^2 \\ &\approx 0.60 \times 10^2 = \$60 \text{ per person} \end{aligned}$$

### Concepts and Vocabulary

- In the expression  $7^2$ , the number 2 is called the exponent.
- In the expression  $-3^7$ , the base is 3.
- The number  $\frac{1}{4^{-2}}$  simplifies to be the positive integer 16.
- The power-of-a-product rule allows us to rewrite  $(5a)^3$  as  $5^3 a^3$ .
- False.  $(-11)^{10} = 11^{10}$
- False. When  $(x^2)^3$  is simplified, the expression becomes  $x^6$ .
- True
- False

**Building Skills**

9.  $3^{-2} = \frac{1}{3^2} = \frac{1}{9}$
10.  $2^{-3} = \frac{1}{2^3} = \frac{1}{8}$
11.  $\left(\frac{1}{2}\right)^{-4} = 2^4 = 16$
12.  $\left(-\frac{1}{2}\right)^{-2} = (-2)^2 = 4$
13.  $7^0 = 1$
14.  $(-8)^0 = 1$
15.  $-(-7)^0 = -1$
16.  $(\sqrt{2})^0 = 1$
17.  $(2^3)^2 = 2^{3 \cdot 2} = 2^6 = 64$
18.  $(3^2)^3 = 3^{2 \cdot 3} = 3^6 = 729$
19.  $(3^2)^{-2} = 3^{2(-2)} = 3^{-4} = \frac{1}{3^4} = \frac{1}{81}$
20.  $(7^2)^{-1} = 7^{2(-1)} = 7^{-2} = \frac{1}{7^2} = \frac{1}{49}$
21.  $(5^{-2})^3 = \left(\frac{1}{5^2}\right)^3 = \left(\frac{1}{25}\right)^3 = \frac{1}{15,625}$
22.  $(5^{-1})^3 = \left(\frac{1}{5}\right)^3 = \left(\frac{1}{5}\right)^3 = \frac{1}{125}$
23.  $(4^{-3}) \cdot (4^5) = 4^{(-3+5)} = 4^2 = 16$
24.  $(7^{-2}) \cdot (7^3) = 7^{(-2+3)} = 7^1 = 7$
25.  $(\sqrt{3})^0 + 10^0 = 1 + 1 = 2$
26.  $(\sqrt{5})^0 - 9^0 = 1 - 1 = 0$
27.  $3^{-2} + \left(\frac{1}{3}\right)^2 = \left(\frac{1}{3}\right)^2 + \left(\frac{1}{3}\right)^2 = \frac{1}{9} + \frac{1}{9} = \frac{2}{9}$
28.  $5^{-2} + \left(\frac{1}{5}\right)^2 = \left(\frac{1}{5}\right)^2 + \left(\frac{1}{5}\right)^2 = \frac{1}{25} + \frac{1}{25} = \frac{2}{25}$
29.  $-2^{-3} = -\frac{1}{2^3} = -\frac{1}{8}$
30.  $-3^{-2} = -\frac{1}{3^2} = -\frac{1}{9}$
31.  $(-3)^{-2} = \frac{1}{(-3)^2} = \frac{1}{9}$
32.  $(-2)^{-3} = \frac{1}{(-2)^3} = -\frac{1}{8}$
33.  $\frac{2^{11}}{2^{10}} = 2^{(11-10)} = 2^1 = 2$
34.  $\frac{3^6}{3^8} = 3^{(6-8)} = 3^{-2} = \frac{1}{3^2} = \frac{1}{9}$
35.  $\frac{(5^3)^4}{5^{12}} = \frac{5^{12}}{5^{12}} = 1$
36.  $\frac{(9^5)^2}{9^8} = \frac{9^{10}}{9^8} = 9^{(10-8)} = 9^2 = 81$
37.  $\frac{2^5 \cdot 3^{-2}}{2^4 \cdot 3^{-3}} = 2^{(5-4)} \cdot 3^{(-2-(-3))} = 2^1 \cdot 3^1 = 6$
38.  $\frac{4^{-2} \cdot 5^3}{4^{-3} \cdot 5} = 4^{(-2-(-3))} \cdot 5^{(3-1)} = 4^1 \cdot 5^2 = 4 \cdot 25 = 100$
39.  $\frac{-5^{-2}}{2^{-1}} = -\frac{1}{5^2} \cdot 2^1 = -\frac{1}{25} \cdot 2 = -\frac{2}{25}$
40.  $\frac{-7^{-2}}{3^{-1}} = -\frac{1}{7^2} \cdot 3^1 = -\frac{1}{49} \cdot 3 = -\frac{3}{49}$
41.  $\left(\frac{2}{3}\right)^{-1} = \frac{3}{2}$
42.  $\left(\frac{1}{5}\right)^{-1} = 5$
43.  $\left(\frac{2}{3}\right)^{-2} = \left(\frac{3}{2}\right)^2 = \frac{3^2}{2^2} = \frac{9}{4}$
44.  $\left(\frac{3}{2}\right)^{-2} = \left(\frac{2}{3}\right)^2 = \frac{2^2}{3^2} = \frac{4}{9}$

$$45. \left(\frac{11}{7}\right)^{-2} = \frac{1}{\left(\frac{11}{7}\right)^2} = \frac{1}{\frac{121}{49}} = \frac{49}{121}$$

$$46. \left(\frac{13}{5}\right)^{-2} = \frac{1}{\left(\frac{13}{5}\right)^2} = \frac{1}{\frac{169}{25}} = \frac{25}{169}$$

$$47. x^4 y^0 = x^4 \cdot 1 = x^4$$

$$48. x^{-1} y^0 = \frac{1}{x} \cdot 1 = \frac{1}{x}$$

$$49. x^{-1} y = \frac{1}{x} \cdot y = \frac{y}{x}$$

$$50. x^2 y^{-2} = x^2 \cdot \frac{1}{y^2} = \frac{x^2}{y^2}$$

$$51. x^{-1} y^{-2} = \frac{1}{xy^2} \quad 52. x^{-3} y^{-2} = \frac{1}{x^3 y^2}$$

$$53. (x^{-3})^4 = x^{-12} = \frac{1}{x^{12}}$$

$$54. (x^{-5})^2 = x^{-10} = \frac{1}{x^{10}}$$

$$55. (x^{-11})^{-3} = x^{(-11) \cdot (-3)} = x^{33}$$

$$56. (x^{-4})^{-12} = x^{(-4) \cdot (-12)} = x^{48}$$

$$57. -3(xy)^5 = -3x^5 y^5$$

$$58. -8(xy)^6 = -8x^6 y^6$$

$$59. 4(xy^{-1})^2 = 4x^2 y^{-2} = \frac{4x^2}{y^2}$$

$$60. 6(x^{-1} y)^3 = 6x^{-3} y^3 = \frac{6y^3}{x^3}$$

$$61. 3(x^{-1} y)^{-5} = 3x^{(-1) \cdot (-5)} y^{-5} = \frac{3x^5}{y^5}$$

$$62. -5(xy^{-1})^{-6} = -5x^{-6} y^{(-1) \cdot (-6)} = -\frac{5y^6}{x^6}$$

$$63. \frac{(x^3)^2}{(x^2)^5} = \frac{x^{3 \cdot 2}}{x^{2 \cdot 5}} = \frac{x^6}{x^{10}} = x^{6-10} = x^{-4} = \frac{1}{x^4}$$

$$64. \frac{x^2}{(x^3)^4} = \frac{x^2}{x^{3 \cdot 4}} = \frac{x^2}{x^{12}} = x^{2-12} = x^{-10} = \frac{1}{x^{10}}$$

$$65. \left(\frac{2xy}{x^2}\right)^3 = \frac{2^3 x^3 y^3}{x^{2 \cdot 3}} = \frac{8x^3 y^3}{x^6} = 8x^{3-6} y^3 = 8x^{-3} y^3 = \frac{8y^3}{x^3}$$

$$66. \left(\frac{5xy}{x^3}\right)^4 = \frac{5^4 x^4 y^4}{x^{3 \cdot 4}} = \frac{625x^4 y^4}{x^{12}} = 625x^{4-12} y^4 = 625x^{-8} y^4 = \frac{625y^4}{x^8}$$

$$67. \left(\frac{-3x^2 y}{x}\right)^5 = \frac{(-3)^5 x^{2 \cdot 5} y^5}{x^5} = \frac{-243x^{10} y^5}{x^5} = -243x^{10-5} y^5 = -243x^5 y^5$$

$$68. \left(\frac{-2xy^2}{y}\right)^3 = \frac{(-2)^3 x^3 y^{2 \cdot 3}}{y^3} = \frac{-8x^3 y^6}{y^3} = -8x^3 y^{6-3} = -8x^3 y^3$$

$$69. \left(\frac{-3x}{5}\right)^{-2} = \frac{(-3)^{-2} x^{-2}}{5^{-2}} = \frac{\frac{1}{9} \cdot \frac{1}{x^2}}{\frac{1}{25}} = \frac{\frac{1}{9x^2}}{\frac{1}{25}} = \frac{25}{9x^2}$$

$$70. \left(\frac{-5y}{3}\right)^{-4} = \frac{(-5)^{-4} y^{-4}}{3^{-4}} = \frac{\frac{1}{5^4} \cdot \frac{1}{y^4}}{\frac{1}{3^4}} = \frac{\frac{1}{625y^4}}{\frac{1}{81}} = \frac{81}{625y^4}$$

$$71. \left(\frac{4x^{-2}}{xy^5}\right)^3 = \frac{4^3 x^{-2 \cdot 3}}{x^3 y^{5 \cdot 3}} = \frac{64x^{-6}}{x^3 y^{15}} = \frac{64}{x^9 y^{15}}$$

$$72. \left(\frac{3x^2 y}{y^3}\right)^5 = \frac{3^5 x^{2 \cdot 5} y^5}{y^{3 \cdot 5}} = \frac{243x^{10} y^5}{y^{15}} = \frac{243x^{10}}{y^{10}}$$

$$73. \frac{x^3 y^{-3}}{x^{-2} y} = x^{3-(-2)} y^{-3-1} = x^5 y^{-4} = \frac{x^5}{y^4}$$

$$74. \frac{x^2 y^{-2}}{x^{-1} y^2} = x^{2-(-1)} y^{-2-2} = x^3 y^{-4} = \frac{x^3}{y^4}$$

$$75. \frac{27x^{-3} y^5}{9x^{-4} y^7} = 3x^{-3-(-4)} y^{5-7} = 3x^1 y^{-2} = \frac{3x}{y^2}$$

$$76. \frac{15x^5y^{-2}}{3x^7y^{-3}} = 5x^{5-7}y^{-2-(-3)} = 5x^{-2}y^1 = \frac{5y}{x^2}$$

$$77. \frac{1}{x^3}(x^2)^3x^{-4} = x^{-3}(x^{2 \cdot 3})x^{-4} = x^{-3}(x^6)x^{-4} \\ = x^{-3+6-4} = x^{-1} = \frac{1}{x}$$

$$78. (8a^3b)^{-4}\left(\frac{2a}{b}\right)^{12} = 8^{-4}a^{3(-4)}b^{-4} \cdot \frac{2^{12}a^{12}}{b^{12}} \\ = (2^3)^{-4}a^{-12}b^{-4} \cdot 2^{12}a^{12}b^{-12} \\ = 2^{3(-4)}a^{-12}b^{-4} \cdot 2^{12}a^{12}b^{-12} \\ = 2^{-12} \cdot 2^{12} \cdot a^{-12} \cdot a^{12} \cdot b^{-4} \cdot b^{-12} \\ = 2^{-12+12} \cdot a^{-12+12} \cdot b^{-4-12} \\ = 2^0a^0b^{-16} = b^{-16} = \frac{1}{b^{16}}$$

$$79. (-xy^2)^3(-2x^2y^2)^{-4} = \frac{(-xy^2)^3}{(-2x^2y^2)^4} \\ = \frac{(-x)^3(y^2)^3}{(-2)^4(x^2)^4(y^2)^4} \\ = \frac{-x^3 \cdot y^{2 \cdot 3}}{16x^{2 \cdot 4}y^{2 \cdot 4}} = \frac{-x^3y^6}{16x^8y^8} \\ = -\frac{1}{16}(x^{3-8})(y^{6-8}) \\ = -\frac{1}{16}x^{-5}y^{-2} = -\frac{1}{16x^5y^2}$$

$$80. \left[\frac{(-x^2y)^3y^{-4}}{(xy)^5}\right]^{-2} = \left[\frac{(-x^{2 \cdot 3})y^3y^{-4}}{x^5y^5}\right]^{-2} \\ = \left(\frac{-x^6y^{-1}}{x^5y^5}\right)^{-2} \\ = (-x^{6-5}y^{-1-5})^{-2} \\ = (-xy^{-6})^{-2} = (-x)^{-2}(y^{-6})^{-2} \\ = x^{-2}y^{(-6)(-2)} = x^{-2}y^{12} = \frac{y^{12}}{x^2}$$

$$81. (4x^3y^2z)^2(x^3y^2z)^{-7} = \frac{(4x^3y^2z)^2}{(x^3y^2z)^7} \\ = \frac{4^2(x^3)^2(y^2)^2z^2}{(x^3)^7(y^2)^7z^7} \\ = \frac{16x^{3 \cdot 2}y^{2 \cdot 2}z^2}{x^{3 \cdot 7}y^{2 \cdot 7}z^7} \\ = \frac{16x^6y^4z^2}{x^{21}y^{14}z^7} \\ = 16x^{6-21}y^{4-14}z^{2-7} \\ = 16x^{-15}y^{-10}z^{-5} \\ = \frac{16}{x^{15}y^{10}z^5}$$

$$82. \frac{(2xyz)^2}{(x^3y)^2(xz)^{-1}} = \frac{2^2x^2y^2z^2}{x^{3 \cdot 2}y^2x^{-1}z^{-1}} = \frac{4x^2y^2z^2}{x^6y^2x^{-1}z^{-1}} \\ = \frac{4x^2y^2z^2}{x^5y^2z^{-1}} = 4x^{2-5}y^{2-2}z^{2-(-1)} \\ = 4x^{-3}y^0z^3 = \frac{4z^3}{x^3}$$

$$83. \frac{5a^{-2}bc^2}{a^4b^{-3}c^2} = 5a^{-2-4}b^{1-(-3)}c^{2-2} \\ = 5a^{-6}b^4 \cdot 1 = \frac{5b^4}{a^6}$$

$$84. \frac{(-3)^2a^5(bc)^2}{(-2)^3a^2b^3c^4} = \frac{9a^5b^2c^2}{-8a^2b^3c^4} = -\frac{9}{8}a^{5-2}b^{2-3}c^{2-4} \\ = -\frac{9}{8}a^3b^{-1}c^{-2} = -\frac{9a^3}{8bc^2}$$

$$85. \left(\frac{xy^{-3}z^{-2}}{x^2y^{-4}z^3}\right)^{-3} = \frac{x^{-3}y^{(-3)(-3)}z^{(-2)(-3)}}{x^{2(-3)}y^{(-4)(-3)}z^{3(-3)}} \\ = \frac{x^{-3}y^9z^6}{x^{-6}y^{12}z^{-9}} \\ = x^{-3-(-6)}y^{9-12}z^{6-(-9)} \\ = x^3y^{-3}z^{15} = \frac{x^3z^{15}}{y^3}$$

$$\begin{aligned}
 86. \quad \left( \frac{xy^{-2}z^{-1}}{x^{-5}yz^{-8}} \right)^{-1} &= \frac{x^{-1}y^{(-2)(-1)}z^{(-1)(-1)}}{x^{(-5)(-1)}y^{-1}z^{(-8)(-1)}} \\
 &= \frac{x^{-1}y^2z^1}{x^5y^{-1}z^8} = x^{-1-5}y^{2-(-1)}z^{1-8} \\
 &= x^{-6}y^3z^{-7} = \frac{y^3}{x^6z^7}
 \end{aligned}$$

87.  $125 = 1.25 \times 10^2$

88.  $247 = 2.47 \times 10^2$

89.  $850,000 = 8.5 \times 10^5$

90.  $205,000 = 2.05 \times 10^5$

91.  $0.007 = 7.0 \times 10^{-3}$

92.  $0.0019 = 1.9 \times 10^{-3}$

93.  $0.00000275 = 2.75 \times 10^{-6}$

94.  $0.0000038 = 3.8 \times 10^{-6}$

**Applying the Concepts**

95.  $135 \text{ ft}^3 \times 2^3 = 1080 \text{ ft}^3$

96.  $675 \text{ in.}^3 \times \left(\frac{1}{3}\right)^3 = 25 \text{ in.}^3$

97. a.  $(2x)^2 = 4x^2 = 4A = 2^2 A$

b.  $(3x)^2 = 9x^2 = 9A = 3^2 A$

98. a.  $\pi \left(\frac{2d}{2}\right)^2 = \pi d^2 = 4\pi \left(\frac{d}{2}\right)^2 = 2^2 A$

b.  $\pi \left(\frac{3d}{2}\right)^2 = 9\pi \left(\frac{d}{2}\right)^2 = 9\pi \left(\frac{d}{2}\right)^2 = 3^2 A$

99.  $F = Sw^2 = 25,000(0.25)^2 = 1562.5 \text{ lb}$

$$\begin{aligned}
 100. \quad F &= S\pi \left(\frac{d}{2}\right)^2 = 10,000(3.14) \left(\frac{1.5}{2}\right)^2 \\
 &\approx 17,662.5 \text{ lbs.}
 \end{aligned}$$

$$\begin{aligned}
 101. \quad 99 \text{ billion} &= 99,000,000,000 = 9.9 \times 10^{10} \\
 330 \text{ million} &= 330,000,000 = 3.3 \times 10^8 \\
 \frac{99 \text{ billion}}{330 \text{ million}} &= \frac{9.9 \times 10^{10}}{3.3 \times 10^8} = \frac{9.9}{3.3} \times \frac{10^{10}}{10^8} \\
 &= 3 \times 10^2 = 300
 \end{aligned}$$

The cost is about \$300 per person.

$$\begin{aligned}
 102. \quad 56 \text{ billion} &= 56,000,000,000 = 5.6 \times 10^{10} \\
 330 \text{ million} &= 330,000,000 = 3.3 \times 10^8 \\
 \frac{56 \text{ billion}}{330 \text{ million}} &= \frac{5.6 \times 10^{10}}{3.3 \times 10^8} = \frac{5.6}{3.3} \times \frac{10^{10}}{10^8} \\
 &\approx 1.7 \times 10^2 = 170
 \end{aligned}$$

The cost is about \$170 per person.

$$\begin{aligned}
 103. \quad 365.25 \text{ days} &\times \frac{24 \text{ hr}}{\text{day}} \times \frac{60 \text{ min}}{\text{hr}} \times \frac{60 \text{ sec}}{\text{min}} \\
 &= 365.25 \times 24 \times 60 \times 60 \text{ sec} \\
 &= 31,557,600 \text{ sec} \\
 &= 3.15576 \times 10^7 \text{ sec}
 \end{aligned}$$

$$\begin{aligned}
 104. \quad 366 \text{ days} &\times \frac{24 \text{ hr}}{\text{day}} \times \frac{60 \text{ min}}{\text{hr}} \times \frac{60 \text{ sec}}{\text{min}} \\
 &= 366 \times 24 \times 60 \times 60 \text{ sec} \\
 &= 31,622,400 \text{ sec} \\
 &= 3.16224 \times 10^7 \text{ sec}
 \end{aligned}$$

| Celestial Body | Equatorial Diameter (km) | Scientific Notation           |
|----------------|--------------------------|-------------------------------|
| Earth          | 12,700                   | $1.27 \times 10^4 \text{ km}$ |
| Moon           | 3480                     | $3.48 \times 10^3 \text{ km}$ |
| Sun            | 1,390,000                | $1.39 \times 10^6 \text{ km}$ |
| Jupiter        | 134,000                  | $1.34 \times 10^5 \text{ km}$ |
| Mercury        | 4800                     | $4.8 \times 10^3 \text{ km}$  |

$$\begin{aligned}
 106. \quad 37,600,000,000,000,000,000,000,000 &= \\
 &= 3.76 \times 10^{22} \text{ atoms}
 \end{aligned}$$

$$\begin{aligned}
 107. \quad 602,000,000,000,000,000,000,000 &= \\
 &= 6.02 \times 10^{20} \text{ atoms}
 \end{aligned}$$

$$\begin{aligned}
 108. \quad 0.000000000000000000000000266 &= \\
 &= 2.66 \times 10^{-20} \text{ kg}
 \end{aligned}$$

$$\begin{aligned}
 109. \quad 0.000000000000000000000000167 &= \\
 &= 1.67 \times 10^{-21} \text{ kg}
 \end{aligned}$$

110.  $380,000,000 = 3.8 \times 10^8 \text{ m}$

$$\begin{aligned}
 111. \quad 5,980,000,000,000,000,000,000,000 &= \\
 &= 5.98 \times 10^{24} \text{ kg}
 \end{aligned}$$

$$\begin{aligned}
 112. \quad 2,000,000,000,000,000,000,000,000,000 &= \\
 &= 2.0 \times 10^{30} \text{ kg}
 \end{aligned}$$

## Beyond the Basics

113.  $2^x = 32$

a.  $2^{x+2} = 2^x \cdot 2^2 = 32 \cdot 4 = 128$

b.  $2^{x-1} = 2^x \cdot 2^{-1} = 32 \cdot \frac{1}{2} = 16$

114.  $3^x = 81$

a.  $3^{x+1} = 3^x \cdot 3^1 = 81 \cdot 3 = 243$

b.  $3^{x-2} = 3^x \cdot 3^{-2} = 81 \cdot \left(\frac{1}{3}\right)^2 = 81 \cdot \frac{1}{9} = 9$

115.  $5^x = 11$

a.  $5^{x+1} = 5^x \cdot 5^1 = 11 \cdot 5 = 55$

b.  $5^{x-2} = 5^x \cdot 5^{-2} = 11 \cdot \left(\frac{1}{5}\right)^2 = 11 \cdot \frac{1}{25} = \frac{11}{25}$

116.  $a^x = b$

a.  $a^{x+2} = a^x \cdot a^2 = b \cdot a^2 = a^2b$

b.  $a^{x-1} = a^x \cdot a^{-1} = b \cdot \frac{1}{a} = \frac{b}{a}$

$$\begin{aligned}
 117. \quad \frac{3^{2-n} \cdot 9^{2n-2}}{3^{3n}} &= \frac{3^{2-n} \cdot (3^2)^{2n-2}}{3^{3n}} \\
 &= \frac{3^{2-n} \cdot 3^{2(2n-2)}}{3^{3n}} \\
 &= \frac{3^{3n-2}}{3^{3n}} = 3^{(3n-2)-3n} \\
 &= 3^{-2} = \frac{1}{3^2} = \frac{1}{9}
 \end{aligned}$$

$$\begin{aligned}
 118. \quad \frac{2^{m+1} \cdot 3^{2m-n} \cdot 5^{m+n+2} \cdot 6^n}{6^m \cdot 10^{n+1} \cdot 15^m} \\
 &= \frac{2^m \cdot 2^1 \cdot 3^{2m} \cdot 3^{-n} \cdot 5^m \cdot 5^n \cdot 5^2 \cdot (2 \cdot 3)^n}{2^m \cdot 2^1 \cdot 3^{2m} \cdot 3^{-n} \cdot 5^m \cdot 5^n \cdot 5^2 \cdot 2^n \cdot 3^n} \\
 &= \frac{2^m \cdot 3^m \cdot 2^{n+1} \cdot 5^{n+1} \cdot 3^m \cdot 5^m}{2^m \cdot 3^m \cdot 2^{n+1} \cdot 5^{n+1} \cdot 3^m \cdot 5^m} \\
 &= \frac{2^{\cancel{m}} \cdot 2^{\cancel{1}} \cdot 3^{2m} \cdot 3^{-n} \cdot 5^{\cancel{m}} \cdot 5^{\cancel{n}} \cdot 5^2 \cdot 2^{\cancel{n}} \cdot 3^{\cancel{n}}}{2^{\cancel{m}} \cdot 3^m \cdot 2^{\cancel{1}} \cdot 2^{\cancel{n}} \cdot 5^{\cancel{m}} \cdot 5^1 \cdot 3^m \cdot 5^{\cancel{m}}} \\
 &= \frac{3^{2m} \cdot 5^2}{3^m \cdot 3^m \cdot 5^1} = \frac{3^{2m} \cdot 5^2}{3^{2m} \cdot 5^1} = 5
 \end{aligned}$$

$$\begin{aligned}
 119. \quad \frac{2^{x(y-z)}}{2^{y(x-z)}} \div \left(\frac{2^y}{2^x}\right)^z &= \frac{2^{xy-xz}}{2^{xy-yz}} \div \frac{2^{yz}}{2^{xz}} \\
 &= \frac{2^{xy}}{2^{xz}} \cdot \frac{2^{xz}}{2^{xy}} \\
 &= \frac{2^{yz}}{2^{xz}} \cdot \frac{2^{xz}}{2^{yz}} = 1
 \end{aligned}$$

$$\begin{aligned}
 120. \quad \left(\frac{a^x}{a^y}\right)^{\frac{1}{xy}} \cdot \left(\frac{a^y}{a^z}\right)^{\frac{1}{yz}} \cdot \left(\frac{a^z}{a^x}\right)^{\frac{1}{xz}} \\
 &= \frac{a^{1/y}}{a^{1/x}} \cdot \frac{a^{1/z}}{a^{1/y}} \cdot \frac{a^{1/x}}{a^{1/z}} = 1
 \end{aligned}$$

## Getting Ready for the Next Section

GR1. a.  $x^2 \cdot x^5 = x^{2+5} = x^7$

b.  $(2x)(-5x^2) = -10x^3$

c.  $(2y^2)(3y^3)(4y^5) = 24y^{2+3+5} = 24y^{10}$

GR2. a.  $2x^2 + 5x^2 = 7x^2$

b.  $3x^2 - 4x^2 = -x^2$

c.  $3x^3 - 5x^3 + 11x^3 = 9x^3$

GR3.  $x(2x+5) = x(2x) + x(5) = 2x^2 + 5x$

GR4.  $2x(7-3x) = 2x(7) + 2x(-3x) = 14x - 6x^2$

$$\begin{aligned}
 \text{GR5. } -5x(x^2+1) &= (-5x)(x^2) + (-5x)(1) \\
 &= -5x^3 - 5x
 \end{aligned}$$

GR6. True. This is an example of the commutative property of multiplication.

GR7. False.  $5x^2$  and  $3x^3$  are not like terms, so they cannot be combined.

$$\begin{aligned}
 \text{GR8. } 2x^2(5x^3-3x+4) &= 10x^{2+3} - 6x^{2+1} + 8x^2 \\
 &= 10x^5 - 6x^3 + 8x^2
 \end{aligned}$$

## P.3 Polynomials

## Practice Problems

1.  $16(7)^2 + 15(7) = 889 \text{ ft}$

2. Using the horizontal method:

$$\begin{aligned}
 & (7x^3 + 2x^2 - 5) + (-2x^3 + 3x^2 + 2x + 1) \\
 &= (7x^3 - 2x^3) + (2x^2 + 3x^2) + 2x + (-5 + 1) \\
 &= 5x^3 + 5x^2 + 2x - 4
 \end{aligned}$$

3. To use the column method, first change the sign in each term in the second polynomial and then add.

$$\begin{array}{r}
 (3x^4 - 5x^3 + 2x^2 + 7) - (-2x^4 + 3x^2 + x - 5) \\
 \begin{array}{r}
 3x^4 \quad - 5x^3 \quad + 2x^2 \quad \quad + 7 \\
 + 2x^4 \quad \quad - 3x^2 \quad - x \quad + 5 \\
 \hline
 5x^4 \quad - 5x^3 \quad - x^2 \quad - x \quad + 12
 \end{array} \\
 (3x^4 - 5x^3 + 2x^2 + 7) - (-2x^4 + 3x^2 + x - 5) \\
 = 5x^4 - 5x^3 - x^2 - x + 12
 \end{array}$$

4.  $-2x^3(4x^2 + 2x - 5)$   
 $= -2x^3(4x^2) - 2x^3(2x) - 2x^3(-5)$   
 $= -8x^5 - 4x^4 + 10x^3$
5.  $(5x^2 + 2x)(-2x^2 + x - 7)$   
 $= 5x^2(-2x^2 + x - 7) + 2x(-2x^2 + x - 7)$   
 $= -10x^4 + 5x^3 - 35x^2 - 4x^3 + 2x^2 - 14x$   
 $= -10x^4 + x^3 - 33x^2 - 14x$
6. a.  $(4x - 1)(x + 7) = 4x^2 + 28x - x - 7$   
 $= 4x^2 + 27x - 7$
- b.  $(3x - 2)(2x - 5) = 6x^2 - 15x - 4x + 10$   
 $= 6x^2 - 19x + 10$
7.  $(3x + 2)^2 = (3x)^2 + 2(3x)(2) + 2^2$   
 $= 9x^2 + 12x + 4$
8.  $(1 - 2x)(1 + 2x) = 1^2 - (2x)^2 = 1 - 4x^2$
9. a.  $(x + 2y)(x - 2y) = x^2 - (2y)^2 = x^2 - 4y^2$
- b.  $(2x - y)^3$   
 $= (2x)^3 - 3(2x)^2y + 3(2x)y^2 - y^3$   
 $= 8x^3 - 3(4x^2)y + 6xy^2 - y^3$   
 $= 8x^3 - 12x^2y + 6xy^2 - y^3$

### Concepts and Vocabulary

1. The polynomial
- $-3x^7 + 2x^2 - 9x + 4$
- has leading coefficient
- $\underline{-3}$
- and degree
- $\underline{7}$
- .

2. When a polynomial is written so that the exponents in each term decrease from left to right, it is said to be in standard form.
3. When a polynomial in  $x$  of degree 3 is added to a polynomial in  $x$  of degree 4, the resulting polynomial has degree  $\underline{4}$ .
4. When a polynomial in  $x$  of degree 3 is multiplied by a polynomial in  $x$  of degree 4, the resulting polynomial has degree  $\underline{7}$ .
5. True
6. True. This is true if  $A$  or  $B$  or both are zero.
7. False. This is not a polynomial because the term  $-3\sqrt{x}$  does not have an exponent that is either a positive integer or zero.
8. True

### Building Skills

9. Polynomial;  $x^2 + 2x + 1$
10. Not a polynomial
11. Not a polynomial
12. A polynomial;  $x^7 + 3x^5 + 3x^4 - 2x + 1$
13. Not a polynomial
14. Not a polynomial
15. A polynomial; in standard form
16. A polynomial; in standard form
17. Degree: 1; terms:  $7x, 3$
18. Degree: 2; terms:  $-3x^2, 7$
19. Degree: 4; terms:  $-x^4, x^2, 2x, -9$
20. Degree: 7; terms:  $9x^7, 2x^3, x, -21$
21.  $(x^3 + 2x^2 - 5x + 3) + (-x^3 + 2x - 4)$   
 $= (x^3 + (-x^3)) + 2x^2 + (-5x + 2x) + (3 - 4)$   
 $= 2x^2 - 3x - 1$
22.  $(x^3 - 3x + 1) + (x^3 - x^2 + x - 3)$   
 $= (x^3 + x^3) - x^2 + (-3x + x) + (1 - 3)$   
 $= 2x^3 - x^2 - 2x - 2$
23.  $(2x^3 - x^2 + x - 5) - (x^3 - 4x + 3)$   
 $= (2x^3 - x^3) - x^2 + (x - (-4x)) + (-5 - 3)$   
 $= x^3 - x^2 + 5x - 8$

$$\begin{aligned} 24. \quad & (-x^3 + 2x - 4) - (x^3 + 3x^2 - 7x + 2) \\ &= (-x^3 - x^3) - 3x^2 + (2x - (-7x)) + (-4 - 2) \\ &= -2x^3 - 3x^2 + 9x - 6 \end{aligned}$$

$$\begin{aligned} 25. \quad & (-2x^4 + 3x^2 - 7x) - (8x^4 + 6x^3 - 9x^2 - 17) \\ &= (-2x^4 - 8x^4) - 6x^3 - 7x + 9x^2 + 17 \\ &= -10x^4 - 6x^3 + 12x^2 - 7x + 17 \end{aligned}$$

$$\begin{aligned} 26. \quad & 3(x^2 - 2x + 2) + 2(5x^2 - x + 4) \\ &= 3x^2 - 6x + 6 + 10x^2 - 2x + 8 \\ &= 13x^2 - 8x + 14 \end{aligned}$$

$$\begin{aligned} 27. \quad & -2(3x^2 + x + 1) + 6(-3x^2 - 2x - 2) \\ &= -6x^2 - 2x - 2 - 18x^2 - 12x - 12 \\ &= -24x^2 - 14x - 14 \end{aligned}$$

$$\begin{aligned} 28. \quad & 2(5x^2 - x + 3) - 4(3x^2 + 7x + 1) \\ &= 10x^2 - 2x + 6 - 12x^2 - 28x - 4 \\ &= -2x^2 - 30x + 2 \end{aligned}$$

$$\begin{aligned} 29. \quad & (3y^3 - 4y + 2) + (2y + 1) - (y^3 - y^2 + 4) \\ &= 3y^3 - 4y + 2 + 2y + 1 - y^3 + y^2 - 4 \\ &= 2y^3 + y^2 - 2y - 1 \end{aligned}$$

$$\begin{aligned} 30. \quad & (5y^2 + 3y - 1) - (y^2 - 2y + 3) + (2y^2 + y + 5) \\ &= 5y^2 + 3y - 1 - y^2 + 2y - 3 + 2y^2 + y + 5 \\ &= 6y^2 + 6y + 1 \end{aligned}$$

$$31. \quad 6x(2x + 3) = 12x^2 + 18x$$

$$32. \quad 7x(3x - 4) = 21x^2 - 28x$$

$$\begin{aligned} 33. \quad & (x + 1)(x^2 + 2x + 2) \\ &= x(x^2 + 2x + 2) + 1(x^2 + 2x + 2) \\ &= x^3 + 2x^2 + 2x + x^2 + 2x + 2 \\ &= x^3 + 3x^2 + 4x + 2 \end{aligned}$$

$$\begin{aligned} 34. \quad & (x - 5)(2x^2 - 3x + 1) \\ &= x(2x^2 - 3x + 1) - 5(2x^2 - 3x + 1) \\ &= 2x^3 - 3x^2 + x - 10x^2 + 15x - 5 \\ &= 2x^3 - 13x^2 + 16x - 5 \end{aligned}$$

$$\begin{aligned} 35. \quad & (3x - 2)(x^2 - x + 1) \\ &= 3x(x^2 - x + 1) - 2(x^2 - x + 1) \\ &= 3x^3 - 3x^2 + 3x - 2x^2 + 2x - 2 \\ &= 3x^3 - 5x^2 + 5x - 2 \end{aligned}$$

$$\begin{aligned} 36. \quad & (2x + 1)(x^2 - 3x + 4) \\ &= 2x(x^2 - 3x + 4) + 1(x^2 - 3x + 4) \\ &= 2x^3 - 6x^2 + 8x + x^2 - 3x + 4 \\ &= 2x^3 - 5x^2 + 5x + 4 \end{aligned}$$

$$\begin{aligned} 37. \quad & (x + 1)(x + 2) = x(x + 2) + 1(x + 2) \\ &= x^2 + 2x + x + 2 = x^2 + 3x + 2 \end{aligned}$$

$$\begin{aligned} 38. \quad & (x + 2)(x + 3) = x(x + 3) + 2(x + 3) \\ &= x^2 + 3x + 2x + 6 \\ &= x^2 + 5x + 6 \end{aligned}$$

$$\begin{aligned} 39. \quad & (3x + 2)(3x + 1) = 9x^2 + 3x + 6x + 2 \\ &= 9x^2 + 9x + 2 \end{aligned}$$

$$\begin{aligned} 40. \quad & (x + 3)(2x + 5) = 2x^2 + 5x + 6x + 15 \\ &= 2x^2 + 11x + 15 \end{aligned}$$

$$\begin{aligned} 41. \quad & (-4x + 5)(x + 3) = -4x^2 - 12x + 5x + 15 \\ &= -4x^2 - 7x + 15 \end{aligned}$$

$$\begin{aligned} 42. \quad & (-2x + 1)(x - 5) = -2x^2 + 10x + x - 5 \\ &= -2x^2 + 11x - 5 \end{aligned}$$

$$\begin{aligned} 43. \quad & (3x - 2)(2x - 1) = 6x^2 - 3x - 4x + 2 \\ &= 6x^2 - 7x + 2 \end{aligned}$$

$$\begin{aligned} 44. \quad & (x - 1)(5x - 3) = 5x^2 - 3x - 5x + 3 \\ &= 5x^2 - 8x + 3 \end{aligned}$$

$$\begin{aligned} 45. \quad & (2x - 3a)(2x + 5a) = 4x^2 + 10ax - 6ax - 15a^2 \\ &= 4x^2 + 4ax - 15a^2 \end{aligned}$$

$$\begin{aligned} 46. \quad & (5x - 2a)(x + 5a) = 5x^2 + 25ax - 2ax - 10a^2 \\ &= 5x^2 + 23ax - 10a^2 \end{aligned}$$

$$47. \quad (x + 2)^2 - x^2 = x^2 + 4x + 4 - x^2 = 4x + 4$$

$$48. \quad (x - 3)^2 - x^2 = x^2 - 6x + 9 - x^2 = -6x + 9$$

$$49. \quad (4x + 1)^2 = 16x^2 + 8x + 1$$

$$50. \quad (3x + 2)^2 = 9x^2 + 12x + 4$$

$$\begin{aligned} 51. \left(x + \frac{3}{4}\right)^2 &= x^2 + 2\left(\frac{3}{4}\right)x + \left(\frac{3}{4}\right)^2 \\ &= x^2 + \frac{3}{2}x + \frac{9}{16} \end{aligned}$$

$$\begin{aligned} 52. \left(x + \frac{2}{5}\right)^2 &= x^2 + 2\left(\frac{2}{5}\right)x + \left(\frac{2}{5}\right)^2 \\ &= x^2 + \frac{4}{5}x + \frac{4}{25} \end{aligned}$$

$$\begin{aligned} 53. (3x - 4y)^2 &= (3x)^2 - 2(3x)(4y) + (4y)^2 \\ &= 9x^2 - 24xy + 16y^2 \end{aligned}$$

$$\begin{aligned} 54. (2x + 5y)^2 &= (2x)^2 + 2(2x)(5y) + (5y)^2 \\ &= 4x^2 + 20xy + 25y^2 \end{aligned}$$

$$\begin{aligned} 55. (x^2 + 2)^2 &= (x^2)^2 + 2(x^2)(2) + 2^2 \\ &= x^4 + 4x^2 + 4 \end{aligned}$$

$$\begin{aligned} 56. (3 - x^2)^2 &= 3^2 - 2(3)(x^2) + (x^2)^2 \\ &= 9 - 6x^2 + x^4 = x^4 - 6x^2 + 9 \end{aligned}$$

$$\begin{aligned} 57. \left(\frac{x}{2} + \frac{2}{x}\right)^2 &= \left(\frac{x}{2}\right)^2 + 2\left(\frac{x}{2}\right)\left(\frac{2}{x}\right) + \left(\frac{2}{x}\right)^2 \\ &= \frac{x^2}{4} + 2 + \frac{4}{x^2} = \frac{x^2}{4} + \frac{4}{x^2} + 2 \end{aligned}$$

$$\begin{aligned} 58. \left(\frac{3}{y} - \frac{y}{3}\right)^2 &= \left(\frac{3}{y}\right)^2 - 2\left(\frac{3}{y}\right)\left(\frac{y}{3}\right) + \left(\frac{y}{3}\right)^2 \\ &= \frac{9}{y^2} - 2 + \frac{y^2}{9} = \frac{9}{y^2} + \frac{y^2}{9} - 2 \end{aligned}$$

$$\begin{aligned} 59. (x + 2)^3 &= x^3 + 3x^2(2) + 3x(2)^2 + 2^3 \\ &= x^3 + 6x^2 + 12x + 8 \end{aligned}$$

$$\begin{aligned} 60. (2x - 1)^3 &= (2x)^3 - 3(2x)^2(1) + 3(2x)(1)^2 - 1^3 \\ &= 8x^3 - 12x^2 + 6x - 1 \end{aligned}$$

$$\begin{aligned} 61. (x + 3)^3 - x^3 &= x^3 + 3x^2(3) + 3x(3)^2 + 3^3 - x^3 \\ &= 9x^2 + 27x + 27 \end{aligned}$$

$$\begin{aligned} 62. (x - 2)^3 - x^3 &= x^3 - 3x^2(2) + 3x(2)^2 - 2^3 - x^3 \\ &= -6x^2 + 12x - 8 \end{aligned}$$

$$\begin{aligned} 63. (x + 2y)^3 &= x^3 + 3x^2(2y) + 3x(2y)^2 + 2^3 \\ &= x^3 + 6x^2y + 12xy^2 + 8 \end{aligned}$$

$$\begin{aligned} 64. (2x + 3y)^3 &= (2x)^3 + 3(2x)^2(3y) + 3(2x)(3y)^2 + (3y)^3 \\ &= 8x^3 + 36x^2y + 54xy^2 + 27y^3 \end{aligned}$$

$$65. (5 + 2x)(5 - 2x) = 5^2 - (2x)^2 = 25 - 4x^2$$

$$66. (3 - 4x)(3 + 4x) = 3^2 - (4x)^2 = 9 - 16x^2$$

$$\begin{aligned} 67. (2x + 3y)(2x - 3y) &= (2x)^2 - (3y)^2 \\ &= 4x^2 - 9y^2 \end{aligned}$$

$$\begin{aligned} 68. (5x - 2y)(5x + 2y) &= (5x)^2 - (2y)^2 \\ &= 25x^2 - 4y^2 \end{aligned}$$

$$69. \left(x + \frac{1}{x}\right)\left(x - \frac{1}{x}\right) = x^2 - \left(\frac{1}{x}\right)^2 = x^2 - \frac{1}{x^2}$$

$$\begin{aligned} 70. \left(\frac{y}{2} - \frac{2}{y}\right)\left(\frac{y}{2} + \frac{2}{y}\right) &= \left(\frac{y}{2}\right)^2 - \left(\frac{2}{y}\right)^2 \\ &= \frac{y^2}{4} - \frac{4}{y^2} \end{aligned}$$

$$71. (x^2 + 3)(x^2 - 3) = (x^2)^2 - 3^2 = x^4 - 9$$

$$72. (x^3 - 2)(x^3 + 2) = (x^3)^2 - 2^2 = x^6 - 4$$

$$\begin{aligned} 73. (2x - 3)(x^2 - 3x + 5) &= 2x(x^2 - 3x + 5) - 3(x^2 - 3x + 5) \\ &= 2x^3 - 6x^2 + 10x - 3x^2 + 9x - 15 \\ &= 2x^3 - 9x^2 + 19x - 15 \end{aligned}$$

$$\begin{aligned} 74. (x - 2)(x^2 - 4x - 3) &= x(x^2 - 4x - 3) - 2(x^2 - 4x - 3) \\ &= x^3 - 4x^2 - 3x - 2x^2 + 8x + 6 \\ &= x^3 - 6x^2 + 5x + 6 \end{aligned}$$

$$\begin{aligned} 75. (1 + y)(1 - y + y^2) &= 1(1 - y + y^2) + y(1 - y + y^2) \\ &= 1 - y + y^2 + y - y^2 + y^3 = y^3 + 1 \end{aligned}$$

$$\begin{aligned} 76. (y + 4)(y^2 - 4y + 16) &= y(y^2 - 4y + 16) + 4(y^2 - 4y + 16) \\ &= y^3 - 4y^2 + 16y + 4y^2 - 16y + 64 \\ &= y^3 + 64 \end{aligned}$$

$$\begin{aligned}
 77. \quad (x-6)(x^2+6x+36) \\
 &= x(x^2+6x+36) - 6(x^2+6x+36) \\
 &= x^3+6x^2+36x-6x^2-36x-216 \\
 &= x^3-216
 \end{aligned}$$

$$\begin{aligned}
 78. \quad (x-1)(x^2+x+1) \\
 &= x(x^2+x+1) - 1(x^2+x+1) \\
 &= x^3+x^2+x-x^2-x-1 = x^3-1
 \end{aligned}$$

$$\begin{aligned}
 79. \quad (x+2y)(3x+5y) &= 3x^2+5xy+6xy+10y^2 \\
 &= 3x^2+11xy+10y^2
 \end{aligned}$$

$$\begin{aligned}
 80. \quad (2x+y)(7x+2y) &= 14x^2+4xy+7xy+2y^2 \\
 &= 14x^2+11xy+2y^2
 \end{aligned}$$

$$\begin{aligned}
 81. \quad (2x-y)(3x+7y) &= 6x^2+14xy-3xy-7y^2 \\
 &= 6x^2+11xy-7y^2
 \end{aligned}$$

$$\begin{aligned}
 82. \quad (x-3y)(2x+5y) &= 2x^2+5xy-6xy-15y^2 \\
 &= 2x^2-xy-15y^2
 \end{aligned}$$

$$\begin{aligned}
 83. \quad (x-y)^2(x+y)^2 &= (x-y)(x-y)(x+y)(x+y) \\
 &= [(x-y)(x+y)][(x-y)(x+y)] \\
 &= (x^2-y^2)(x^2-y^2) \\
 &= x^4-2x^2y^2+y^4
 \end{aligned}$$

$$\begin{aligned}
 84. \quad (2x+y)^2(2x-y)^2 \\
 &= (2x+y)(2x+y)(2x-y)(2x-y) \\
 &= [(2x-y)(2x+y)][(2x-y)(2x+y)] \\
 &= (4x^2-y^2)(4x^2-y^2) \\
 &= 16x^4-8x^2y^2+y^4
 \end{aligned}$$

$$\begin{aligned}
 85. \quad (x+y)(x-2y)^2 &= (x+y)(x^2-4xy+4y^2) \\
 &= x(x^2-4xy+4y^2) + y(x^2-4xy+4y^2) \\
 &= x^3-4x^2y+4xy^2+x^2y-4xy^2+4y^3 \\
 &= x^3-3x^2y+4y^3
 \end{aligned}$$

$$\begin{aligned}
 86. \quad (x-y)(x+2y)^2 &= (x-y)(x^2+4xy+4y^2) \\
 &= x(x^2+4xy+4y^2) - y(x^2+4xy+4y^2) \\
 &= x^3+4x^2y+4xy^2-x^2y-4xy^2-4y^3 \\
 &= x^3+3x^2y-4y^3
 \end{aligned}$$

$$\begin{aligned}
 87. \quad (x-2y)^3(x+2y) \\
 &= (x^3+3x^2(-2y)+3x(-2y)^2+(-2y)^3) \cdot (x+2y) \\
 &= (x^3-6x^2y+12xy^2-8y^3)(x+2y) \\
 &= x(x^3-6x^2y+12xy^2-8y^3) \\
 &\quad + 2y(x^3-6x^2y+12xy^2-8y^3) \\
 &= x^4-6x^3y+12x^2y^2-8xy^3 \\
 &\quad + 2x^3y-12x^2y^2+24xy^3-16y^4 \\
 &= x^4-4x^3y+16xy^3-16y^4
 \end{aligned}$$

$$\begin{aligned}
 88. \quad (2x+y)^3(2x-y) \\
 &= [(2x)^3+3(2x)^2y+3(2x)y^2+y^3] \cdot (2x-y) \\
 &= (8x^3+12x^2y+6xy^2+y^3)(2x-y) \\
 &= 2x(8x^3+12x^2y+6xy^2+y^3) \\
 &\quad - y(8x^3+12x^2y+6xy^2+y^3) \\
 &= 16x^4+24x^3y+12x^2y^2+2xy^3 \\
 &\quad - 8x^3y-12x^2y^2-6xy^3-y^4 \\
 &= 16x^4+16x^3y-4xy^3-y^4
 \end{aligned}$$

For exercises 89–94, use

$$a^2+b^2=(a+b)^2-2ab=(a-b)^2+2ab.$$

$$\begin{aligned}
 89. \quad x+y &= 4, \quad xy = 3 \\
 x^2+y^2 &= (x+y)^2-2xy \\
 &= 4^2-2 \cdot 3 = 16-6 = 10
 \end{aligned}$$

$$\begin{aligned}
 90. \quad x-y &= 3, \quad xy = 2 \\
 x^2+y^2 &= (x-y)^2+2xy \\
 &= 3^2+2 \cdot 2 = 9+4 = 13
 \end{aligned}$$

$$91. \text{ a. } x + \frac{1}{x} = 3$$

$$\text{Let } a = x \text{ and } b = \frac{1}{x}. \text{ Then } ab = x\left(\frac{1}{x}\right) = 1.$$

$$\begin{aligned}
 x^2 + \frac{1}{x^2} &= \left(x + \frac{1}{x}\right)^2 - 2x\left(\frac{1}{x}\right) \\
 &= 3^2 - 2 = 9 - 2 = 7
 \end{aligned}$$

- b. Let  $a = x^2$  and  $b = \frac{1}{x^2}$ . Using the result

from part a, we have

$$x^4 + \frac{1}{x^4} = \left(x^2 + \frac{1}{x^2}\right)^2 - 2x^2\left(\frac{1}{x^2}\right) \\ = 7^2 - 2 = 49 - 2 = 47$$

92. a.  $x - \frac{1}{x} = 2$

Let  $a = x$  and  $b = \frac{1}{x}$ . Then  $ab = x\left(\frac{1}{x}\right) = 1$ .

$$x^2 + \frac{1}{x^2} = \left(x - \frac{1}{x}\right)^2 + 2x\left(\frac{1}{x}\right) \\ = 2^2 + 2 = 4 + 2 = 6$$

- b. Let  $a = x^2$  and  $b = \frac{1}{x^2}$ . Using the result

from part a, we have

$$x^4 + \frac{1}{x^4} = \left(x^2 + \frac{1}{x^2}\right)^2 - 2x^2\left(\frac{1}{x^2}\right) \\ = 6^2 - 2 = 36 - 2 = 34$$

93. Let  $a = 3x$  and  $b = 2y$ . Then  $ab = 6xy$ .

$$9x^2 + 4y^2 = (3x + 2y)^2 - 2(6xy) \\ = 12^2 - 2 \cdot 6 \cdot 6 = 144 - 72 = 72$$

94. Let  $a = 3x$  and  $b = 7y$ . Then  $ab = 21xy$ .

$$9x^2 + 49y^2 = (3x - 7y)^2 + 2(21xy) \\ = 10^2 + 2(21)(-1) \\ = 100 - 42 = 58$$

### Applying the Concepts

95.  $0.11(5^2) + 0.02(5) + 5.57 = \$8.425$  in 2015  
(five years after 2010)

96.  $0.035(4^2) + 0.15(4) + 5.17 = 6.33$

In 2012 (four years after 2008), theater grosses were 6.33 billion dollars.

97.  $0.1(40^2) + 40 + 50 = \$250.00$

98.  $(15 - 10)^2 + 5(15) = 5^2 + 75 \\ = 25 + 75 = \$100.00$

99.  $d = 16(5^2) + 20(5) = 16(25) + 100 = 500$  feet

100.  $d = 16(2^2) + 10(2) = 16(4) + 20 = 84$  feet

101. a.  $-x + 22.50$

b.  $(30 + 10x)(22.50 - x) \\ = 30(22.50) - 30x + 225x - 10x^2 \\ = 675 + 195x - 10x^2 \\ = -10x^2 + 195x + 675$

102. a.  $10n + 250$

b.  $(50 - 2n)(10n + 250) \\ = 50(10n) + 50(250) - 2n(10n) - 2n(250) \\ = 500n + 12,500 - 20n^2 - 500n \\ = -20n^2 + 12,500$

### Beyond the Basics

103.  $(a + b + c)^2 = (a + b + c)(a + b + c) \\ = a(a + b + c) + b(a + b + c) + c(a + b + c) \\ = a^2 + ab + ac + ab + b^2 + bc + ac + bc + c^2 \\ = a^2 + b^2 + c^2 + 2ab + 2bc + 2ac$

104.  $(x^2 + x + 1)^2 = (x^2)^2 + x^2 + 1^2 + 2x^2x + 2x(1) + 2x^2(1) \\ = x^4 + x^2 + 1 + 2x^3 + 2x + 2x^2 \\ = x^4 + 2x^3 + 3x^2 + 2x + 1$

105.  $(2x + y - z)^2 = (2x)^2 + y^2 + (-z)^2 + 2(2x)y + 2y(-z) + 2(2x)(-z) \\ = 4x^2 + y^2 + z^2 + 4xy - 2yz - 4xz$   
 $(2x - y + z)^2 = (2x)^2 + (-y)^2 + z^2 + 2(2x)(-y) + 2(-y)z + 2(2x)z \\ = 4x^2 + y^2 + z^2 - 4xy - 2yz + 4xz$

(continued on next page)

(continued)

$$\begin{aligned}(2x + y - z)^2 - (2x - y + z)^2 &= (4x^2 + y^2 + z^2 + 4xy - 2yz - 4xz) - (4x^2 + y^2 + z^2 - 4xy - 2yz + 4xz) \\ &= 8xy - 8xz\end{aligned}$$

Alternatively, recognize the difference of two squares.

$$\begin{aligned}(2x + y - z)^2 - (2x - y + z)^2 &= [(2x + y - z) + (2x - y + z)][(2x + y - z) - (2x - y + z)] \\ &= (4x)(2y - 2z) = 8xy - 8xz\end{aligned}$$

**106.**  $a + b + c = 8$ ;  $ab + bc + ac = 12$

$$\begin{aligned}(a + b + c)^2 &= a^2 + b^2 + c^2 + 2ab + 2bc + 2ac = a^2 + b^2 + c^2 + 2(ab + bc + ac) \\ 8^2 &= a^2 + b^2 + c^2 + 2(12) \\ 64 &= a^2 + b^2 + c^2 + 24 \\ 40 &= a^2 + b^2 + c^2\end{aligned}$$

**107.**  $x + y + z = 12$ ;  $x^2 + y^2 + z^2 = 44$

$$\begin{aligned}(x + y + z)^2 &= x^2 + y^2 + z^2 + 2xy + 2yz + 2xz = x^2 + y^2 + z^2 + 2(xy + yz + xz) \\ 12^2 &= 44 + 2(xy + yz + xz) \\ 144 &= 44 + 2(xy + yz + xz) \\ 100 &= 2(xy + yz + xz) \\ 50 &= xy + yz + xz\end{aligned}$$

**108.**  $x^2 + y^2 + z^2 = 64$ ;  $xy + yz + xz = 18$

$$\begin{aligned}(x + y + z)^2 &= x^2 + y^2 + z^2 + 2xy + 2yz + 2xz = x^2 + y^2 + z^2 + 2(xy + yz + xz) \\ (x + y + z)^2 &= 64 + 2(18) \\ (x + y + z)^2 &= 100 \\ x + y + z &= \sqrt{100} = 10\end{aligned}$$

**109.**  $(a + b + c)(a^2 + b^2 + c^2 - ab - bc - ac)$

$$\begin{aligned}&= a(a^2 + b^2 + c^2 - ab - bc - ac) + b(a^2 + b^2 + c^2 - ab - bc - ac) + c(a^2 + b^2 + c^2 - ab - bc - ac) \\ &= a^3 + ab^2 + ac^2 - a^2b - abc - a^2c + a^2b + b^3 + bc^2 - ab^2 - b^2c - abc + a^2c + b^2c + c^3 - abc - bc^2 - ac^2 \\ &= a^3 + b^3 + c^3 - 3abc\end{aligned}$$

**110.** From exercise 109,

$$(0)(a^2 + b^2 + c^2 - ab - bc - ac) = a^3 + b^3 + c^3 - 3abc \Rightarrow 0 = a^3 + b^3 + c^3 - 3abc \Rightarrow a^3 + b^3 + c^3 = 3abc$$

**111.** Let  $a = x - y$ ,  $b = y - z$ , and  $c = z - x$ . Then  $a + b + c = (x - y) + (y - z) + (z - x) = 0$ .From exercise 110, if  $a + b + c = 0$ , then  $a^3 + b^3 + c^3 = 3abc$ . So,

$$(x - y)^3 + (y - z)^3 + (z - x)^3 = 3(x - y)(y - z)(z - x).$$

**112.** Let  $a = 2x - 3y$ ,  $b = 3y - 5z$ , and  $c = 5z - 3x$ . Then  $a + b + c = (2x - 3y) + (3y - 5z) + (5z - 3x) = 0$ .From exercise 110, if  $a + b + c = 0$ , then  $a^3 + b^3 + c^3 = 3abc$ . So,

$$(2x - 3y)^3 + (3y - 5z)^3 + (5z - 3x)^3 = 3(2x - 3y)(3y - 5z)(5z - 3x).$$

113. From exercise 109,  $(a+b+c)(a^2+b^2+c^2-ab-bc-ac)=a^3+b^3+c^3-3abc$ .

If  $a+b+c=8$  and  $ab+bc+ac=19$ , we have

$$(a+b+c)(a^2+b^2+c^2-ab-bc-ac)=(a+b+c)(a^2+b^2+c^2-(ab+bc+ac))=a^3+b^3+c^3-3abc \Rightarrow 8(a^2+b^2+c^2-19)=a^3+b^3+c^3-3abc \quad (1)$$

From exercise 103,

$$(a+b+c)^2=a^2+b^2+c^2+2ab+2bc+2ac=a^2+b^2+c^2+2(ab+bc+ac) \Rightarrow 8^2=a^2+b^2+c^2+2(19) \Rightarrow 64=a^2+b^2+c^2+38 \Rightarrow a^2+b^2+c^2=26$$

Substituting into equation (1) gives us  $8(26-19)=a^3+b^3+c^3-3abc \Rightarrow 56=a^3+b^3+c^3-3abc$ .

114. From exercise 109,  $(a+b+c)(a^2+b^2+c^2-ab-bc-ac)=a^3+b^3+c^3-3abc$ .

If  $a+b+c=9$  and  $ab+bc+ca=11$ , we have

$$(a+b+c)(a^2+b^2+c^2-ab-bc-ac)=(a+b+c)(a^2+b^2+c^2-(ab+bc+ac))=a^3+b^3+c^3-3abc \Rightarrow 9(a^2+b^2+c^2-11)=a^3+b^3+c^3-3abc \quad (1)$$

From exercise 103,

$$(a+b+c)^2=a^2+b^2+c^2+2ab+2bc+2ac=a^2+b^2+c^2+2(ab+bc+ac) \Rightarrow 9^2=a^2+b^2+c^2+2(11) \Rightarrow 81=a^2+b^2+c^2+22 \Rightarrow a^2+b^2+c^2=59$$

Substituting into equation (1) gives us  $9(59-11)=a^3+b^3+c^3-3abc \Rightarrow 432=a^3+b^3+c^3-3abc$ .

### Getting Ready for the Next Section

GR1.  $x(2x+5)=x(2x)+x(5)=2x^2+5x$

GR2.  $2x(7-3x)=2x(7)+2x(-3x)=14x-6x^2$

GR3.  $-5x(x^2+1)=-5x(x^2)-5x(1)=-5x^3-5x$

GR4. True. This is an example of the commutative property for multiplication.

|       | $a$       | $b$       | $a+b$    | $ab$       |
|-------|-----------|-----------|----------|------------|
| GR5.  | 3         | 4         | <u>7</u> | <u>12</u>  |
| GR6.  | -3        | 5         | <u>2</u> | <u>-15</u> |
| GR7.  | <u>4</u>  | 2         | 6        | 8          |
| GR8.  | <u>3</u>  | -5        | -2       | -15        |
| GR9.  | <u>2</u>  | <u>5</u>  | 7        | 10         |
| GR10. | <u>3</u>  | <u>5</u>  | 8        | 15         |
| GR11. | <u>-5</u> | <u>7</u>  | 2        | -35        |
| GR12. | <u>5</u>  | <u>-7</u> | -2       | -35        |
| GR13. | <u>-2</u> | <u>-3</u> | -5       | 6          |
| GR14. | <u>2</u>  | <u>3</u>  | 5        | 6          |

### P.4 Factoring Polynomials

#### Practice Problems

1. a.  $6x^5+14x^3=2x^3(3x^2+7)$

b.  $7x^5+21x^4+35x^2=7x^2(x^3+3x^2+5)$

c.  $5x^2(x-y)+2(x-y)=(x-y)(5x^2+2)$

2. a.  $x^2+6x+8=(x+4)(x+2)$

b.  $x^2-3x-10=(x-5)(x+2)$

3. a.  $x^2+4x+4=(x+2)^2$

b.  $9x^2-6x+1=(3x-1)^2$

4. a.  $x^2-16=(x-4)(x+4)$

b.  $4x^2-25=(2x-5)(2x+5)$

5.  $x^4-81=(x^2-9)(x^2+9)$   
 $= (x-3)(x+3)(x^2+9)$

6. a.  $x^3 - 125 = (x - 5)(x^2 + 5x + 25)$
- b.  $27x^3 + 8 = (3x + 2)(9x^2 - 6x + 4)$
7. Following the reasoning in Example 7, in four years, the company will have invested the initial 12 million dollars, plus an additional 4 million dollars. Thus, the total investment is 16 million dollars. To find the profit or loss with 16 million dollars invested, let  $x = 16$  in the profit-loss polynomial:
- $$(0.012)(x - 14)(x^2 + 14x + 196)$$
- $$= (0.012)(16 - 14)(16^2 + 14 \cdot 16 + 196)$$
- $$= 16.224$$
- The company will have made a profit of 16.224 million dollars in four years.
8. a.  $x^3 + 3x^2 + x + 3 = x^2(x + 3) + 1(x + 3)$   
 $= (x + 3)(x^2 + 1)$
- b.  $28x^3 - 20x^2 - 7x + 5$   
 $= 4x^2(7x - 5) - (7x - 5)$   
 $= (7x - 5)(4x^2 - 1)$   
 $= (7x - 5)(2x + 1)(2x - 1)$
- c.  $x^2 - y^2 - 2y - 1 = x^2 - (y^2 + 2y + 1)$   
 $= x^2 - (y + 1)^2$   
 $= (x - (y + 1))(x + (y + 1))$   
 $= (x - y - 1)(x + y + 1)$
9.  $x^4 + 5x^2 + 9 = x^4 + 6x^2 - x^2 + 9$   
 $= x^4 + 6x^2 + 9 - x^2$   
 $= (x^2 + 3)^2 - x^2$   
 $= [(x^2 + 3) + x][(x^2 + 3) - x]$   
 $= (x^2 + x + 3)(x^2 - x + 3)$
10. a.  $5x^2 + 11x + 2 = 5x^2 + 10x + x + 2$   
 $= (5x^2 + 10x) + (x + 2)$   
 $= 5x(x + 2) + 1(x + 2)$   
 $= (5x + 1)(x + 2)$
- b.  $4x^2 + x - 3 = 4x^2 + 4x - 3x - 3$   
 $= (4x^2 + 4x) - (3x + 3)$   
 $= 4x(x + 1) - 3(x + 1)$   
 $= (4x - 3)(x + 1)$

### Concepts and Vocabulary

- The polynomials  $x + 2$  and  $x - 2$  are called factors of the polynomial  $x^2 - 4$ .
- The polynomial  $3y$  is the greatest common monomial factor of the polynomial  $3y^2 + 6y$ .
- The GCF of the polynomial  $10x^3 + 30x^2$  is  $10x^2$ .
- A polynomial that cannot be factored as a product of two polynomials (excluding constant polynomials  $\pm 1$ ) is said to be irreducible.
- True
- True
- False. The polynomial  $x^2 - 4$  can be factored as  $(x + 2)(x - 2)$ . Therefore, it is not irreducible.
- True

### Building Skills

- $8x - 24 = 8(x - 3)$
- $5x + 25 = 5(x + 5)$
- $-6x^2 + 12x = -6x(x - 2)$
- $-3x^2 + 21 = -3(x^2 - 7)$
- $7x^2 + 14x^3 = 7x^2(1 + 2x)$
- $9x^3 - 18x^4 = 9x^3(1 - 2x)$
- $x^4 + 2x^3 + x^2 = x^2(x^2 + 2x + 1)$
- $x^4 - 5x^3 + 7x^2 = x^2(x^2 - 5x + 7)$
- $3x^3 - x^2 = x^2(3x - 1)$
- $2x^3 + 2x^2 = 2x^2(x + 1)$
- $8ax^3 + 4ax^2 = 4ax^2(2x + 1)$
- $ax^4 - 2ax^2 + ax = ax(x^3 - 2x + 1)$
- $x(x - y) + 3(x - y) = (x - y)(x + 3)$
- $2a(a + 1) - 3(a + 1) = (a + 1)(2a - 3)$

$$23. \quad 3x(2x+y)+5y(2x+y)=(2x+y)(3x+5y)$$

$$24. \quad \begin{aligned} 2x(5x-3y)+7y(5x-3y) \\ = (5x-3y)(2x+7y) \end{aligned}$$

$$25. \quad \begin{aligned} (y+2)^2-3(y+2) &= (y+2)(y+2-3) \\ &= (y+2)(y-1) \end{aligned}$$

$$26. \quad \begin{aligned} (y+1)^2+(y+1) &= (y+1)((y+1)+1) \\ &= (y+1)(y+2) \end{aligned}$$

$$27. \quad \begin{aligned} x^3+3x^2+x+3 &= x^2(x+3)+1(x+3) \\ &= (x+3)(x^2+1) \end{aligned}$$

$$28. \quad \begin{aligned} x^3+5x^2+x+5 &= x^2(x+5)+1(x+5) \\ &= (x+5)(x^2+1) \end{aligned}$$

$$29. \quad \begin{aligned} x^3-5x^2+x-5 &= x^2(x-5)+1(x-5) \\ &= (x-5)(x^2+1) \end{aligned}$$

$$30. \quad \begin{aligned} x^3-7x^2+x-7 &= x^2(x-7)+1(x-7) \\ &= (x-7)(x^2+1) \end{aligned}$$

$$31. \quad \begin{aligned} 6x^3+4x^2+3x+2 &= 2x^2(3x+2)+1(3x+2) \\ &= (3x+2)(2x^2+1) \end{aligned}$$

$$32. \quad \begin{aligned} 3x^3+6x^2+x+2 &= 3x^2(x+2)+1(x+2) \\ &= (x+2)(3x^2+1) \end{aligned}$$

$$33. \quad \begin{aligned} 12x^7+4x^5+3x^4+x^2 \\ &= 4x^5(3x^2+1)+x^2(3x^2+1) \\ &= (3x^2+1)(4x^5+x^2) \\ &= x^2(3x^2+1)(4x^3+1) \end{aligned}$$

$$34. \quad \begin{aligned} 3x^7+3x^5+x^4+x^2 &= 3x^5(x^2+1)+x^2(x^2+1) \\ &= (x^2+1)(3x^5+x^2) \\ &= x^2(x^2+1)(3x^3+1) \end{aligned}$$

$$35. \quad \begin{aligned} xy+ab+bx+ay &= (xy+bx)+(ab+ay) \\ &= x(y+b)+a(b+y) \\ &= x(y+b)+a(y+b) \\ &= (b+y)(x+a) \end{aligned}$$

$$36. \quad \begin{aligned} 2a^2x-b-bx+2a^2 &= 2a^2x+2a^2-(b+bx) \\ &= 2a^2(x+1)-b(1+x) \\ &= 2a^2(x+1)-b(x+1) \\ &= (x+1)(2a^2-b) \end{aligned}$$

$$37. \quad x^2+7x+12=(x+3)(x+4)$$

$$38. \quad x^2+8x+15=(x+3)(x+5)$$

$$39. \quad x^2-6x+8=(x-2)(x-4)$$

$$40. \quad x^2-9x+14=(x-2)(x-7)$$

$$41. \quad x^2-3x-4=(x+1)(x-4)$$

$$42. \quad x^2-5x-6=(x+1)(x-6)$$

$$43. \quad \text{Irreducible} \qquad 44. \quad \text{Irreducible}$$

$$45. \quad 2x^2+x-36=(2x+9)(x-4)$$

$$46. \quad 2x^2+3x-27=(2x+9)(x-3)$$

$$47. \quad 6x^2+17x+12=(2x+3)(3x+4)$$

$$48. \quad 8x^2-10x-3=(2x-3)(4x+1)$$

$$49. \quad 3x^2-11x-4=(3x+1)(x-4)$$

$$50. \quad 5x^2+7x+2=(5x+2)(x+1)$$

$$51. \quad \text{Irreducible} \qquad 52. \quad \text{Irreducible}$$

$$53. \quad x^2-xy-20y^2=(x+4y)(x-5y)$$

$$54. \quad 8p^2-2pq-15q^2=(4p+5q)(2p-3q)$$

$$55. \quad \begin{aligned} 15x^2-28+x &= 15x^2+x-28 \\ &= (3x-4)(5x+7) \end{aligned}$$

$$56. \quad \begin{aligned} 24y+35+4y^2 &= 4y^2+24y+35 \\ &= (2y+7)(2y+5) \end{aligned}$$

$$57. \quad x^2+6x+9=(x+3)^2$$

$$58. \quad x^2+8x+16=(x+4)^2$$

$$59. \quad 9x^2+6x+1=(3x+1)^2$$

$$60. \quad 36x^2+12x+1=(6x+1)^2$$

$$61. \quad 25x^2-20x+4=(5x-2)^2$$

$$\begin{aligned} 62. \quad 64x^2 + 32x + 4 &= 4(16x^2 + 8x + 1) \\ &= 4(4x + 1)^2 \end{aligned}$$

$$63. \quad 49x^2 + 42x + 9 = (7x + 3)^2$$

$$64. \quad 9x^2 + 24x + 16 = (3x + 4)^2$$

$$65. \quad x^2 - 64 = (x - 8)(x + 8)$$

$$66. \quad x^2 - 121 = (x - 11)(x + 11)$$

$$67. \quad 4x^2 - 1 = (2x - 1)(2x + 1)$$

$$68. \quad 9x^2 - 1 = (3x - 1)(3x + 1)$$

$$69. \quad 16x^2 - 9 = (4x - 3)(4x + 3)$$

$$70. \quad 25x^2 - 49 = (5x - 7)(5x + 7)$$

$$71. \quad x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$\begin{aligned} 72. \quad x^4 - 81 &= (x^2 - 9)(x^2 + 9) \\ &= (x - 3)(x + 3)(x^2 + 9) \end{aligned}$$

$$73. \quad 20x^4 - 5 = 5(4x^4 - 1) = 5(2x^2 - 1)(2x^2 + 1)$$

$$74. \quad 12x^4 - 75 = 3(4x^4 - 25) = 3(2x^2 - 5)(2x^2 + 5)$$

$$75. \quad x^3 + 64 = x^3 + 4^3 = (x + 4)(x^2 - 4x + 16)$$

$$76. \quad x^3 + 125 = x^3 + 5^3 = (x + 5)(x^2 - 5x + 25)$$

$$77. \quad x^3 - 27 = x^3 - 3^3 = (x - 3)(x^2 + 3x + 9)$$

$$78. \quad x^3 - 216 = x^3 - 6^3 = (x - 6)(x^2 + 6x + 36)$$

$$79. \quad 8 - x^3 = 2^3 - x^3 = (2 - x)(4 + 2x + x^2)$$

$$80. \quad 27 - x^3 = 3^3 - x^3 = (3 - x)(9 + 3x + x^2)$$

$$81. \quad 8x^3 - 27 = (2x)^3 - 3^3 = (2x - 3)(4x^2 + 6x + 9)$$

$$\begin{aligned} 82. \quad 8x^3 - 125 &= (2x)^3 - 5^3 \\ &= (2x - 5)(4x^2 + 10x + 25) \end{aligned}$$

$$\begin{aligned} 83. \quad 40x^3 + 5 &= 5(8x^3 + 1) = 5((2x)^3 + 1^3) \\ &= 5(2x + 1)(4x^2 - 2x + 1) \end{aligned}$$

$$\begin{aligned} 84. \quad 7x^3 + 56 &= 7(x^3 + 8) = 7(x^3 + 2^3) \\ &= 7(x + 2)(x^2 - 2x + 4) \end{aligned}$$

$$\begin{aligned} 85. \quad x^4 + x^2 + 25 &= x^4 + 10x^2 - 9x^2 + 25 \\ &= (x^4 + 10x^2 + 25) - 9x^2 \\ &= (x^2 + 5)^2 - 9x^2 \\ &= [(x^2 + 5) + 3x][(x^2 + 5) - 3x] \\ &= (x^2 + 3x + 5)(x^2 - 3x + 5) \end{aligned}$$

$$\begin{aligned} 86. \quad x^4 + x^2 + 1 &= x^4 + 2x^2 - x^2 + 1 \\ &= (x^4 + 2x^2 + 1) - x^2 \\ &= (x^2 + 1)^2 - x^2 \\ &= [(x^2 + 1) + x][(x^2 + 1) - x] \\ &= (x^2 + x + 1)(x^2 - x + 1) \end{aligned}$$

$$\begin{aligned} 87. \quad x^4 + 15x^2 + 64 &= x^4 + 16x^2 - x^2 + 64 \\ &= (x^4 + 16x^2 + 64) - x^2 \\ &= (x^2 + 8)^2 - x^2 \\ &= [(x^2 + 8) + x][(x^2 + 8) - x] \\ &= (x^2 + x + 8)(x^2 - x + 8) \end{aligned}$$


$$\begin{aligned} 88. \quad x^4 - 7x^2 + 9 &= x^4 - 6x^2 - x^2 + 9 \\ &= (x^4 - 6x^2 + 9) - x^2 \\ &= (x^2 - 3)^2 - x^2 \\ &= [(x^2 - 3) + x][(x^2 - 3) - x] \\ &= (x^2 + x - 3)(x^2 - x - 3) \end{aligned}$$

$$\begin{aligned} 89. \quad x^4 - x^2 + 16 &= x^4 + 8x^2 - 9x^2 + 16 \\ &= (x^4 + 8x^2 + 16) - 9x^2 \\ &= (x^2 + 4)^2 - 9x^2 \\ &= [(x^2 + 4) + 3x][(x^2 + 4) - 3x] \\ &= (x^2 + 3x + 4)(x^2 - 3x + 4) \end{aligned}$$


90.  $x^4 - 16x^2 + 36 = x^4 - 12x^2 - 4x^2 + 36$   
 $= (x^4 - 12x^2 + 36) - 4x^2$   
 $= (x^2 - 6)^2 - 4x^2$   
 $= [(x^2 - 6) + 2x][(x^2 - 6) - 2x]$   
 $= (x^2 + 2x - 6)(x^2 - 2x - 6)$
91.  $x^4 + 4 = x^4 + 4x^2 - 4x^2 + 4$   
 $= (x^4 + 4x^2 + 4) - 4x^2$   
 $= (x^2 + 2)^2 - 4x^2$   
 $= [(x^2 + 2) + 2x][(x^2 + 2) - 2x]$   
 $= (x^2 + 2x + 2)(x^2 - 2x + 2)$
92.  $x^4 + 64 = x^4 + 16x^2 - 16x^2 + 64$   
 $= (x^4 + 16x^2 + 64) - 16x^2$   
 $= (x^2 + 8)^2 - 16x^2$   
 $= [(x^2 + 8) + 4x][(x^2 + 8) - 4x]$   
 $= (x^2 + 4x + 8)(x^2 - 4x + 8)$
93.  $1 - 16x^2 = (1 + 4x)(1 - 4x)$
94.  $4 - 25x^2 = (2 - 5x)(2 + 5x)$
95.  $x^2 - 6x + 9 = (x - 3)^2$
96.  $x^2 - 8x + 16 = (x - 4)^2$
97.  $4x^2 + 4x + 1 = (2x + 1)^2$
98.  $16x^2 + 8x + 1 = (4x + 1)^2$
99.  $2x^2 - 8x - 10 = 2(x^2 - 4x - 5)$   
 $= 2(x + 1)(x - 5)$
100.  $5x^2 - 10x - 40 = 5(x^2 - 2x - 8)$   
 $= 5(x + 2)(x - 4)$
101.  $2x^2 + 3x - 20 = (2x - 5)(x + 4)$
102.  $2x^2 - 7x - 30 = (2x + 5)(x - 6)$
103.  $x^2 - 24x + 36$  is irreducible.
104.  $x^2 - 20x + 25$  is irreducible.
105.  $3x^5 + 12x^4 + 12x^3 = 3x^3(x^2 + 4x + 4)$   
 $= 3x^3(x + 2)^2$
106.  $2x^5 + 16x^4 + 32x^3 = 2x^3(x^2 + 8x + 16)$   
 $= 2x^3(x + 4)^2$
107.  $9x^2 - 1 = (3x + 1)(3x - 1)$
108.  $16x^2 - 25 = (4x + 5)(4x - 5)$
109.  $16x^2 + 24x + 9 = (4x + 3)^2$
110.  $4x^2 + 20x + 25 = (2x + 5)^2$
111.  $x^2 + 15$  is irreducible.
112.  $x^2 + 24$  is irreducible.
113.  $45x^3 + 8x^2 - 4x = x(45x^2 + 8x - 4)$   
 $= x(5x + 2)(9x - 2)$
114.  $25x^3 + 40x^2 - 9x = x(25x^2 + 40x - 9)$   
 $= x(5x + 9)(5x - 1)$
115.  $ax^2 - 7a^2x - 8a^3 = a(x^2 - 7ax - 8a^2)$   
 $= a(x + a)(x - 8a)$
116.  $ax^2 - 10a^2x - 24a^3 = a(x^2 - 10ax - 24a^2)$   
 $= a(x + 2a)(x - 12a)$
117.  $x^2 - 16a^2 + 8x + 16 = (x^2 + 8x + 16) - 16a^2$   
 $= (x + 4)^2 - 16a^2$   
 $= (x + 4 + 4a)(x + 4 - 4a)$
118.  $x^2 - 36a^2 + 6x + 9 = (x^2 + 6x + 9) - 36a^2$   
 $= (x + 3)^2 - 36a^2$   
 $= (x + 3 + 6a)(x + 3 - 6a)$
119.  $3x^5 + 12x^4y + 12x^3y^2 = 3x^3(x^2 + 4xy + 4y^2)$   
 $= 3x^3(x + 2y)^2$
120.  $4x^5 + 24x^4y + 36x^3y^2 = 4x^3(x^2 + 6xy + 9y^2)$   
 $= 4x^3(x + 3y)^2$
121.  $x^2 - 25a^2 + 4x + 4 = (x^2 + 4x + 4) - 25a^2$   
 $= (x + 2)^2 - 25a^2$   
 $= (x + 2 + 5a)(x + 2 - 5a)$

122.  $x^2 - 16a^2 + 4x + 4 = (x^2 + 4x + 4) - 16a^2$   
 $= (x + 2)^2 - 16a^2$   
 $= (x + 2 + 4a)(x + 2 - 4a)$
123.  $18x^6 + 12x^5y + 2x^4y^2 = 2x^4(9x^2 + 6xy + y^2)$   
 $= 2x^4(3x + y)^2$
124.  $12x^6 + 12x^5y + 3x^4y^2 = 3x^4(4x^2 + 4xy + y^2)$   
 $= 3x^4(2x + y)^2$

### Applying the Concepts



If one side of the garden is  $x$  feet, and the perimeter is 16 feet, then  $\frac{16 - 2x}{2} = 8 - x$  gives the other dimension of the rectangle. So, the area of the garden is  $x(8 - x)$ .



If one side of the tray is  $x$  inches, and the perimeter is 42 inches, then  $\frac{42 - 2x}{2} = 21 - x$  gives the other dimension of the tray. So, the area of the tray is  $x(21 - x)$ .

127. a. Mateo will break even when the profit or loss is 0, so we must determine in how many months that will occur. Evaluate the polynomial that represents Mateo's profit or loss for different values until the value of the polynomial is 0. The number of months is represented by  $x$ .

| $x$ | $10x^3 - 30x^2 - 90x - 50$       | Polynomial value |
|-----|----------------------------------|------------------|
| 0   | $10(0)^3 - 30(0)^2 - 90(0) - 50$ | -50              |
| 1   | $10(1)^3 - 30(1)^2 - 90(1) - 50$ | -160             |

| $x$ | $10x^3 - 30x^2 - 90x - 50$       | Polynomial value |
|-----|----------------------------------|------------------|
| 2   | $10(2)^3 - 30(2)^2 - 90(2) - 50$ | -270             |
| 3   | $10(3)^3 - 30(3)^2 - 90(3) - 50$ | -320             |
| 4   | $10(4)^3 - 30(4)^2 - 90(4) - 50$ | -250             |
| 5   | $10(5)^3 - 30(5)^2 - 90(5) - 50$ | 0                |

Mateo will break even in 5 months.


- b. Evaluate the profit or loss after 10 months by letting  $x = 10$ .  
 $10(10)^3 - 30(10)^2 - 90(10) - 50 = 6050$   
 After 10 months, there is a profit of \$6050.
128. a. The company will break even when the profit or loss is 0, so we must determine in how many months that will occur. Evaluate the polynomial that represents the company's profit or loss for different values until the value of the polynomial is 0. The number of months is represented by  $x$ .

| $x$ | $90x^3 - 360x^2 + 630x - 2520$       | Polynomial value |
|-----|--------------------------------------|------------------|
| 0   | $90(0)^3 - 360(0)^2 + 630(0) - 2520$ | -2520            |
| 1   | $90(1)^3 - 360(1)^2 + 630(1) - 2520$ | -2160            |
| 2   | $90(2)^3 - 360(2)^2 + 630(2) - 2520$ | -1980            |
| 3   | $90(3)^3 - 360(3)^2 + 630(3) - 2520$ | -1440            |
| 4   | $90(4)^3 - 360(4)^2 + 630(4) - 2520$ | 0                |

The company will break even in 4 months.


- b. Evaluate the profit or loss after 10 months by letting  $x = 10$ .  
 $90(10)^3 - 360(10)^2 + 630(10) - 2520 = 57,780$   
 After 10 months, there is a profit of \$57,780.

Use the figure below for exercises 129 and 130.




129. If  $x$  = the length of the cut corner, then  $36 - 2x$  = the length of the box, and  $16 - 2x$  = the width of the box. The height of the box is  $x$ . So,  $v = x(36 - 2x)(16 - 2x) = x(2(18 - x))(2(8 - x)) = 4x(18 - x)(8 - x)$ .
130.  $S.A. = (36 - 2x)(16 - 2x) + 2x(36 - 2x) + 2x(16 - 2x) = 576 - 104x + 4x^2 + 72x - 4x^2 + 32x - 4x^2 = 576 - 4x^2 = 4(144 - x^2) = 4(12 - x)(12 + x)$
131. The area of the outside circle =  $4\pi \text{ cm}^2$ . The area of the inside circle =  $\pi x^2 \text{ cm}^2$ . So, the area of the disk = area of outside circle – area of inside circle =  $4\pi - \pi x^2 = \pi(4 - x^2) = \pi(2 - x)(2 + x) \text{ cm}^2$ .

132.



The volume of the inside cylinder is  $3^2\pi \cdot 8 = 72\pi \text{ ft}^3$ , and the volume of the outside cylinder is  $x^2\pi \cdot 8 = 8\pi x^2 \text{ ft}^3$ .  
So the volume between the cylinders is  
 $8\pi x^2 - 72\pi = 8\pi(x^2 - 9) = 8\pi(x - 3)(x + 3) \text{ ft}^3$ .

133.



If one side of the fence is  $x$  feet, and the rancher needs a total of 2800 feet of fencing, then the width of the fence is  $2800 - 2x$ . So, the area of the pen is  $x(2800 - 2x) = 2800x - 2x^2 = 2x(1400 - x) \text{ ft}^2$ .

134. The area of the figure = the area of the rectangle plus the area of the circle. Find the length of the rectangle as follows: The circumference of the circle =  $2\pi\left(\frac{x}{2}\right) = \pi x$ . The perimeter of the figure is 48, so the length

of the rectangle is  $\frac{48 - \pi x}{2}$ . The area of the circle is  $\frac{\pi x^2}{4}$  and the area of the rectangle is

$$x\left(24 - \frac{\pi x}{2}\right) = 24x - \frac{\pi x^2}{2}. \text{ So, the area of the figure is}$$

$$\frac{\pi x^2}{4} + \left(24x - \frac{\pi x^2}{2}\right) = \frac{1}{4}(96x - \pi x^2) = \frac{1}{4}x(96 - \pi x) \text{ in.}^2$$

### Beyond the Basics

$$\begin{aligned} 135. \quad x^3 + y^3 + x + y &= (x + y)(x^2 - xy + y^2) + x + y \\ &= (x + y)(x^2 - xy + y^2 + 1) \end{aligned}$$

$$\begin{aligned} 136. \quad x^4 - 11x^2 + 1 &= x^4 - 2x^2 + 1 - 9x^2 = (x^2 - 1)^2 - 9x^2 \\ &= (x^2 - 1 - 3x)(x^2 - 1 + 3x) = (x^2 - 3x - 1)(x^2 + 3x - 1) \end{aligned}$$

$$\begin{aligned}
 137. \quad x^4 + y^4 + x^2y^2 &= x^4 + 2x^2y^2 + y^4 - x^2y^2 = (x^2 + y^2)^2 - x^2y^2 \\
 &= (x^2 + y^2 - xy)(x^2 + y^2 + xy) = (x^2 - xy + y^2)(x^2 + xy + y^2)
 \end{aligned}$$

$$\begin{aligned}
 138. \quad x^4 + 4y^4 &= (x^2)^2 + (2y^2)^2 + 4x^2y^2 - 4x^2y^2 \\
 &= (x^2)^2 + 4x^2y^2 + (2y^2)^2 - 4x^2y^2 \\
 &= (x^2 + 2y^2)^2 - 4x^2y^2 \\
 &= (x^2 + 2y^2 - 2xy)(x^2 + 2y^2 + 2xy) \\
 &= (x^2 - 2xy + 2y^2)(x^2 + 2xy + 2y^2)
 \end{aligned}$$

$$\begin{aligned}
 139. \quad x^6 - 64 &= (x^3)^2 - 8^2 = (x^3 - 8)(x^3 + 8) \\
 &= (x - 2)(x + 2x + 4)(x + 2)(x - 2x + 4) \\
 &= (x - 2)(x + 2)(x + 2x + 4)(x - 2x + 4)
 \end{aligned}$$

$$140. \quad x^6 - 26x^3 - 27 = (x^3 - 27)(x^3 + 1) = (x - 3)(x^2 + 3x + 9)(x + 1)(x^2 - x + 1)$$

$$\begin{aligned}
 141. \quad (x^2 - 3x)^2 - 38(x^2 - 3x) - 80 &= [(x^2 - 3x) - 40][(x^2 - 3x) + 2] \\
 &= (x^2 - 3x - 40)(x^2 - 3x + 2) \\
 &= (x - 8)(x + 5)(x - 2)(x - 1)
 \end{aligned}$$

$$\begin{aligned}
 142. \quad 4x^2 - y^2 - z^2 + 2yz &= 4x^2 - (y^2 + z^2 - 2yz) = 4x^2 - (y - z)^2 \\
 &= 4x^2 - (y - z)^2 = (2x - (y - z))(2x + (y - z)) \\
 &= (2x - y + z)(2x + y - z)
 \end{aligned}$$

$$143. \quad \text{From exercise 109 in Section P.3, we have } (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ac) = a^3 + b^3 + c^3 - 3abc.$$

$$\begin{aligned}
 \text{Then, } a^3 + b^3 + c^3 - 3abc &= (a + b + c)(a^2 + b^2 + c^2 - ab - bc - ac) \\
 &= \frac{1}{2}(a + b + c) \cdot 2(a^2 + b^2 + c^2 - ab - bc - ac) \\
 &= \frac{1}{2}(a + b + c)(2a^2 + 2b^2 + 2c^2 - 2ab - 2bc - 2ac) \\
 &= \frac{1}{2}(a + b + c)((a^2 - 2ab + b^2) + (b^2 - 2bc + c^2) + (c^2 - 2ca + a^2)) \\
 &= \frac{1}{2}(a + b + c)[(a - b)^2 + (b - c)^2 + (c - a)^2]
 \end{aligned}$$

If  $a^3 + b^3 + c^3 - 3abc = 0$ , then either  $a + b + c = 0$  or  $(a - b)^2 + (b - c)^2 + (c - a)^2 = 0 \Rightarrow a = b = c$ .

$$144. \quad \text{See exercise 111 in section P.3.}$$

$$(x - y)^3 + (y - z)^3 + (z - x)^3 = 3(x - y)(y - z)(z - x)$$

$$145. \quad \text{Let } a = z(x - y) = xz - yz, \quad b = x(y - z) = xy - xz, \quad \text{and } c = y(z - x) = yz - xy. \text{ Then}$$

$a + b + c = xz - yz + xy - xz + yz - xy = 0$ , and we can apply Section P.3 exercise 110.

$$z^3(x - y)^3 + x^3(y - z)^3 + y^3(z - x)^3 = 3xyz(x - y)(y - z)(z - x)$$

$$146. \frac{(x^2 - y^2)^3 + (y^2 - z^2)^3 + (z^2 - x^2)^3}{(x - y)^3 + (y - z)^3 + (z - x)^3}$$

In the numerator, let  $a = x^2 - y^2$ ,  $b = y^2 - z^2$ , and  $c = z^2 - x^2$ . Then  $a + b + c = 0$ , and we can apply Section P.3 exercise 111. Use the result from exercise 144 above for the denominator.

$$\begin{aligned} \frac{(x^2 - y^2)^3 + (y^2 - z^2)^3 + (z^2 - x^2)^3}{(x - y)^3 + (y - z)^3 + (z - x)^3} &= \frac{3(x^2 - y^2)(y^2 - z^2)(z^2 - x^2)}{3(x - y)(y - z)(z - x)} \\ &= \frac{(x + y)(x - y)(y + z)(y - z)(z + x)(z - x)}{(x - y)(y - z)(z - x)} \\ &= (x + y)(y + z)(z + x) \end{aligned}$$

### Getting Ready for the Next Section

$$\text{GR1. } 3x^2 + 9 = 3(x^2 + 3)$$

$$\text{GR2. } 21x - 7x^2 = 7x(3 - x)$$

$$\text{GR3. } x^2 + 2x - 15 = (x + 5)(x - 3)$$

$$\text{GR4. } x^2 - 5x + 4 = (x - 4)(x - 1)$$

$$\text{GR5. } \frac{2x}{4x + 6} = \frac{2x}{2(2x + 3)} = \frac{x}{2x + 3}$$

$$\text{GR6. } \frac{2x^3 + 3x}{x^2 + x} = \frac{x(2x^2 + 3)}{x(x + 1)} = \frac{2x^2 + 3}{x + 1}$$

$$\text{GR7. } \frac{x^2 + 3x + 2}{x^2 + 4x + 3} = \frac{(x + 1)(x + 2)}{(x + 1)(x + 3)} = \frac{x + 2}{x + 3}$$

$$\text{GR8. } \frac{x^2 - x - 6}{x^2 - 4} = \frac{(x - 3)(x + 2)}{(x - 2)(x + 2)} = \frac{x - 3}{x - 2}$$

## P.5 Rational Expressions

### Practice Problems

$$1. \frac{0.05(1 - 0.1)}{0.95(0.1) + (0.05)(1 - 0.1)} \approx 0.32$$

The likelihood that a student who tests positive is a nonuser when the test that is used is 95% accurate is about 32%.

$$2. \text{ a. Note that } x \neq 0, x \neq -4.$$

$$\frac{2x^3 + 8x^2}{3x^2 + 12x} = \frac{2x^2(x + 4)}{3x(x + 4)} = \frac{2x}{3}$$

$$\text{b. Note that } x \neq -2.$$

$$\begin{aligned} \frac{x^2 - 4}{x^2 + 4x + 4} &= \frac{(x - 2)(x + 2)}{(x + 2)^2} \\ &= \frac{(x - 2)\cancel{(x + 2)}}{(x + 2)\cancel{(x + 2)}} = \frac{x - 2}{x + 2} \end{aligned}$$

$$3. \text{ Note that } x \neq 0, x \neq -4, x \neq -1$$

$$\begin{aligned} \frac{x^2 - 2x - 3}{\frac{7x^3 + 28x^2}{4x + 4}} &= \frac{x^2 - 2x - 3}{7x^3 + 28x^2} \div \frac{4x + 4}{2x^4 + 8x^3} \\ &= \frac{x^2 - 2x - 3}{7x^3 + 28x^2} \cdot \frac{2x^4 + 8x^3}{4x + 4} \\ &= \frac{(x - 3)\cancel{(x + 1)} \cdot \cancel{2}x^3 \cdot \cancel{(x + 4)}}{7x^2 \cdot \cancel{(x + 4)} \cdot \cancel{4} \cdot \cancel{(x + 1)}} \\ &= \frac{x(x - 3)}{14} \end{aligned}$$

$$4. \text{ a. Note that } x \neq 6, x \neq -6.$$

$$\begin{aligned} \frac{5x + 22}{x^2 - 36} + \frac{2(x + 10)}{x^2 - 36} &= \frac{5x + 22 + 2x + 20}{x^2 - 36} \\ &= \frac{7x + 42}{x^2 - 36} \\ &= \frac{7\cancel{(x + 6)}}{(x - 6)\cancel{(x + 6)}} = \frac{7}{x - 6} \end{aligned}$$

$$\text{b. Note that } x \neq -4, x \neq 3.$$

$$\begin{aligned} \frac{4x + 1}{x^2 + x - 12} - \frac{3x + 4}{x^2 + x - 12} &= \frac{(4x + 1) - (3x + 4)}{x^2 + x - 12} \\ &= \frac{x - 3}{x^2 + x - 12} \\ &= \frac{\cancel{x - 3}}{(x + 4)\cancel{(x - 3)}} \\ &= \frac{1}{x + 4} \end{aligned}$$

5. a. Note that  $x \neq -2$ ,  $x \neq 5$ .

$$\begin{aligned} \frac{2x}{x+2} + \frac{3x}{x-5} &= \frac{2x(x-5)}{(x+2)(x-5)} + \frac{3x(x+2)}{(x+2)(x-5)} \\ &= \frac{2x^2 - 10x + 3x^2 + 6x}{(x+2)(x-5)} = \frac{5x^2 - 4x}{(x+2)(x-5)} \\ &= \frac{x(5x-4)}{(x+2)(x-5)} \end{aligned}$$

- b. Note that  $x \neq 4$ ,  $x \neq -3$ .

$$\begin{aligned} \frac{5x}{x-4} - \frac{2x}{x+3} &= \frac{5x(x+3)}{(x-4)(x+3)} - \frac{2x(x-4)}{(x-4)(x+3)} \\ &= \frac{5x^2 + 15x - 2x^2 + 8x}{(x-4)(x+3)} = \frac{3x^2 + 23x}{(x-4)(x+3)} \\ &= \frac{x(3x+23)}{(x-4)(x+3)} \end{aligned}$$

6. a.  $\frac{x^2+3x}{x^2(x+2)^2(x-2)}, \frac{4x^2+1}{3x(x-2)^2}$   
 LCD =  $3x^2(x+2)^2(x-2)^2$

b.  $\frac{2x-1}{x^2-25}, \frac{3-7x^2}{x^2+4x-5}$   
 $x^2-25 = (x-5)(x+5)$   
 $x^2+4x-5 = (x-1)(x+5)$   
 LCD =  $(x-1)(x-5)(x+5)$

7. a. Note that  $x \neq 2$ ,  $x \neq -2$ .

$$\begin{aligned} \frac{4}{x^2-4x+4} + \frac{x}{x^2-4} &= \frac{4}{(x-2)^2} + \frac{x}{(x-2)(x+2)} \quad \text{The LCD is } (x-2)^2(x+2). \\ &= \frac{4(x+2)}{(x-2)^2(x+2)} + \frac{x(x-2)}{(x-2)^2(x+2)} \\ &= \frac{(4x+8) + (x^2-2x)}{(x-2)^2(x+2)} = \frac{x^2+2x+8}{(x-2)^2(x+2)} \end{aligned}$$

- b. Note that  $x \neq 0$ ,  $x \neq 5$ .

$$\begin{aligned} \frac{2x}{3(x-5)^2} - \frac{6x}{2(x^2-5x)} &= \frac{2x}{3(x-5)^2} - \frac{6x}{2x(x-5)} \quad \text{The LCD is } 6x(x-5)^2. \\ &= \frac{2x(2x)}{3 \cdot 2x(x-5)^2} - \frac{6x(3)(x-5)}{3 \cdot 2x(x-5)^2} \\ &= \frac{4x^2 - 18x^2 + 90x}{6x(x-5)^2} = \frac{-14x^2 + 90x}{6x(x-5)^2} \\ &= \frac{2x(-7x+45)}{6x(x-5)^2} = \frac{-7x+45}{3(x-5)^2} \end{aligned}$$

8. Note that  $x \neq 0$ ,  $x \neq 5$ ,  $x \neq -5$ .

$$\begin{aligned} \frac{\frac{5}{3x} + \frac{1}{3}}{\frac{x^2-25}{3x}} &= \frac{\left(\frac{5}{3x} + \frac{1}{3}\right) \cdot 3x}{\left(\frac{x^2-25}{3x}\right) \cdot 3x} = \frac{5+x}{x^2-25} \\ &= \frac{\cancel{x+5}}{(x-5)\cancel{(x+5)}} = \frac{1}{x-5} \end{aligned}$$

9. Note that  $x \neq \frac{4}{5}$ .

$$\begin{aligned} \frac{5x}{3x - \frac{4}{2 - \frac{1}{3}}} &= \frac{5x}{3x - \frac{4}{\frac{5}{3}}} = \frac{5x}{3x - 4\left(\frac{3}{5}\right)} = \frac{5x}{3x - \frac{12}{5}} \\ &= \frac{5x \cdot 5}{\left(3x - \frac{12}{5}\right) \cdot 5} = \frac{25x}{15x - 12} \end{aligned}$$

### Concepts and Vocabulary

1. The least common denominator of two rational expressions is the polynomial of least degree that contains each denominator as a factor.
2. The first step in finding the LCD of two rational expressions is to factor the denominators completely.
3. If the denominators of two rational expressions are  $x^2 - 2x$  and  $x^2 - x - 2$ , then the LCD is  $x(x-2)(x+1)$ .
4. A rational expression that contains another rational expression in its numerator or denominator is called a complex rational expression.

5. False 6. True

7. False 8. True

**Building Skills**

$$9. \frac{2x+2}{x^2+2x+1} = \frac{2(x+1)}{(x+1)^2} = \frac{2}{x+1}, x \neq -1$$

$$10. \frac{3x-6}{x^2-4x+4} = \frac{3(x-2)}{(x-2)^2} = \frac{3}{x-2}, x \neq 2$$

$$11. \frac{3x+3}{x^2-1} = \frac{3(x+1)}{(x-1)(x+1)} = \frac{3}{x-1}, x \neq -1, x \neq 1$$

$$12. \frac{10-5x}{4-x^2} = \frac{5(2-x)}{(2-x)(2+x)} = \frac{5}{2+x}, x \neq -2, x \neq 2$$

$$13. \frac{2x-6}{9-x^2} = \frac{2(x-3)}{(3-x)(3+x)} = \frac{-2(3-x)}{(3-x)(3+x)} \\ = -\frac{2}{3+x}, x \neq -3, x \neq 3$$

$$14. \frac{15+3x}{x^2-25} = \frac{3(5+x)}{(x-5)(x+5)} = \frac{3}{x-5}, x \neq -5, x \neq 5$$

$$15. \frac{2x-1}{1-2x} = \frac{-1(1-2x)}{1-2x} = -1, x \neq \frac{1}{2}$$

$$16. \frac{2-5x}{5x-2} = \frac{-1(5x-2)}{5x-2} = -1, x \neq \frac{2}{5}$$

$$17. \frac{x^2-6x+9}{4x-12} = \frac{(x-3)^2}{4(x-3)} = \frac{x-3}{4}, x \neq 3$$

$$18. \frac{x^2-10x+25}{3x-15} = \frac{(x-5)^2}{3(x-5)} = \frac{x-5}{3}, x \neq 5$$

$$19. \frac{7x^2+7x}{x^2+2x+1} = \frac{7x(x+1)}{(x+1)^2} = \frac{7x}{x+1}, x \neq -1$$

$$20. \frac{4x^2+12x}{x^2+6x+9} = \frac{4x(x+3)}{(x+3)^2} = \frac{4x}{x+3}, x \neq -3$$

$$21. \frac{x^2-11x+10}{x^2+6x-7} = \frac{(x-10)(x-1)}{(x+7)(x-1)} \\ = \frac{x-10}{x+7}, x \neq -7, x \neq 1$$

$$22. \frac{x^2+2x-15}{x^2-7x+12} = \frac{(x+5)(x-3)}{(x-4)(x-3)} \\ = \frac{x+5}{x-4}, x \neq 3, x \neq 4$$

$$23. \frac{6x^4+14x^3+4x^2}{6x^4-10x^3-4x^2} = \frac{2x^2(3x^2+7x+2)}{2x^2(3x^2-5x-2)} \\ = \frac{2x^2(3x+1)(x+2)}{2x^2(3x+1)(x-2)} \\ = \frac{x+2}{x-2}, x \neq -\frac{1}{3}, x \neq 0, x \neq 2$$

$$24. \frac{3x^3+x^2}{3x^4-11x^3-4x^2} = \frac{x^2(3x+1)}{x^2(3x^2-11x-4)} \\ = \frac{x^2(3x+1)}{x^2(3x+1)(x-4)} \\ = \frac{1}{x-4}, x \neq -\frac{1}{3}, x \neq 0, x \neq 4$$

$$25. \frac{x-3}{2x+4} \cdot \frac{10x+20}{5x-15} = \frac{x-3}{2(x+2)} \cdot \frac{10(x+2)}{5(x-3)} = 1$$

$$26. \frac{6x+4}{2x-8} \cdot \frac{x-4}{9x+6} = \frac{2(3x+2)}{2(x-4)} \cdot \frac{x-4}{3(3x+2)} = \frac{1}{3}$$

$$27. \frac{2x+6}{4x-8} \cdot \frac{x^2+x-6}{x^2-9} = \frac{2(x+3)}{4(x-2)} \cdot \frac{(x+3)(x-2)}{(x+3)(x-3)} \\ = \frac{x+3}{2(x-3)}$$

$$28. \frac{25x^2-9}{4-2x} \cdot \frac{4-x^2}{10x-6} \\ = \frac{(5x+3)(5x-3)}{2(2-x)} \cdot \frac{(2-x)(2+x)}{2(5x-3)} \\ = \frac{(5x+3)(x+2)}{4}$$

$$29. \frac{x^2-7x}{x^2-6x-7} \cdot \frac{x^2-1}{x^2} \\ = \frac{x(x-7)}{(x+1)(x-7)} \cdot \frac{(x+1)(x-1)}{x^2} = \frac{x-1}{x}$$

$$30. \frac{x^2-9}{x^2-6x+9} \cdot \frac{5x-15}{x+3} = \frac{(x-3)(x+3)}{(x-3)(x-3)} \cdot \frac{5(x-3)}{x+3} \\ = 5$$

$$31. \frac{x^2-x-6}{x^2+3x+2} \cdot \frac{x^2-1}{x^2-9} \\ = \frac{(x-3)(x+2)}{(x+1)(x+2)} \cdot \frac{(x-1)(x+1)}{(x-3)(x+3)} = \frac{x-1}{x+3}$$

$$32. \frac{x^2+2x-8}{x^2+x-20} \cdot \frac{x^2-16}{x^2+5x+4} = \frac{(x-2)(x+4)}{(x-4)(x+5)} \cdot \frac{(x-4)(x+4)}{(x+1)(x+4)} = \frac{(x-2)(x+4)}{(x+5)(x+1)}$$

$$33. \frac{2-x}{x+1} \cdot \frac{x^2+3x+2}{x^2-4} = \frac{-1(x-2)}{x+1} \cdot \frac{(x+1)(x+2)}{(x-2)(x+2)} = -1$$

$$34. \frac{3-x}{x+5} \cdot \frac{x^2+8x+15}{x^2-9} = \frac{-1(x-3)}{x+5} \cdot \frac{(x+3)(x+5)}{(x-3)(x+3)} = -1$$

$$35. \frac{x+2}{6} \div \frac{4x+8}{9} = \frac{x+2}{6} \cdot \frac{9}{4(x+2)} = \frac{3}{8}$$

$$36. \frac{x+3}{20} \div \frac{4x+12}{9} = \frac{x+3}{20} \cdot \frac{9}{4(x+3)} = \frac{9}{80}$$

$$37. \frac{x^2-9}{x} \div \frac{2x+6}{5x^2} = \frac{(x-3)(x+3)}{x} \cdot \frac{5x^2}{2(x+3)} = \frac{5x(x-3)}{2}$$

$$38. \frac{x^2-1}{3x} \div \frac{7x-7}{x^2+x} = \frac{(x-1)(x+1)}{3x} \cdot \frac{x(x+1)}{7(x-1)} = \frac{(x+1)^2}{21}$$

$$39. \frac{x^2+2x-3}{x^2+8x+16} \div \frac{x-1}{3x+12} = \frac{(x-1)(x+3)}{(x+4)^2} \cdot \frac{3(x+4)}{x-1} = \frac{3(x+3)}{x+4}$$

$$40. \frac{x^2+5x+6}{x^2+6x+9} \div \frac{x^2+3x+2}{x^2+7x+12} = \frac{(x+2)(x+3)}{(x+3)(x+3)} \cdot \frac{(x+3)(x+4)}{(x+1)(x+2)} = \frac{x+4}{x+1}$$

$$41. \left( \frac{x^2-9}{x^3+8} \div \frac{x+3}{x^3+2x^2-x-2} \right) \left( \frac{1}{x^2-1} \right) = \frac{(x-3)(x+3)}{(x+2)(x^2-2x+4)} \cdot \frac{(x^2-1)(x+2)}{x+3} \cdot \frac{1}{x^2-1} = \frac{x-3}{x^2-2x+4}$$

$$42. \left( \frac{x^2-25}{x^2-3x-4} \div \frac{x^2+3x-10}{x^2-1} \right) \left( \frac{x-2}{x-5} \right) = \frac{(x-5)(x+5)}{(x-4)(x+1)} \cdot \frac{(x-1)(x+1)}{(x+5)(x-2)} \cdot \frac{x-2}{x-5} = \frac{x-1}{x-4}$$

$$43. \frac{x}{5} + \frac{3}{5} = \frac{x+3}{5}$$

$$44. \frac{7}{4} - \frac{x}{4} = \frac{7-x}{4}$$

$$45. \frac{x}{2x+1} + \frac{4}{2x+1} = \frac{x+4}{2x+1}$$

$$46. \frac{2x}{7x-3} + \frac{x}{7x-3} = \frac{3x}{7x-3}$$

$$47. \frac{x^2}{x+1} - \frac{x^2-1}{x+1} = \frac{x^2-(x^2-1)}{x+1} = \frac{1}{x+1}$$

$$48. \frac{2x+7}{3x+2} - \frac{x-2}{3x+2} = \frac{(2x+7)-(x-2)}{3x+2} = \frac{x+9}{3x+2}$$

$$49. \frac{4}{3-x} + \frac{2x}{x-3} = \frac{4}{-1(x-3)} + \frac{2x}{x-3} \\ = \frac{-4+2x}{x-3} = \frac{2(x-2)}{x-3}$$

$$50. \frac{-2}{1-x} + \frac{2-x}{x-1} = \frac{-2}{1-x} + \frac{2-x}{-1(1-x)} \\ = \frac{-2}{1-x} - \frac{2-x}{1-x} = \frac{-4+x}{1-x} \\ = \frac{x-4}{1-x} \text{ or } -\frac{x-4}{x-1}$$

$$51. \frac{5x}{x^2+1} + \frac{2x}{x^2+1} = \frac{7x}{x^2+1}$$

$$52. \frac{x}{2(x-1)^2} + \frac{3x}{2(x-1)^2} = \frac{4x}{2(x-1)^2} = \frac{2x}{(x-1)^2}$$

$$53. \frac{7x}{2(x-3)} + \frac{x}{2(x-3)} = \frac{8x}{2(x-3)} = \frac{4x}{x-3}$$

$$54. \frac{4x}{4(x+5)^2} + \frac{8x}{4(x+5)^2} = \frac{12x}{4(x+5)^2} = \frac{3x}{(x+5)^2}$$

$$55. \frac{x}{x^2-4} - \frac{2}{x^2-4} = \frac{x-2}{(x-2)(x+2)} = \frac{1}{x+2}$$

$$56. \frac{5x}{x^2-1} - \frac{5}{x^2-1} = \frac{5x-5}{(x-1)(x+1)} = \frac{5(x-1)}{(x-1)(x+1)} = \frac{5}{x+1}$$

$$57. \frac{x-2}{2x+1} - \frac{x}{2x-1} = \frac{(x-2)(2x-1)}{(2x+1)(2x-1)} - \frac{x(2x+1)}{(2x+1)(2x-1)} = \frac{(2x^2-5x+2)-(2x^2+x)}{(2x+1)(2x-1)}$$

$$= \frac{-6x+2}{(2x+1)(2x-1)} = \frac{2(1-3x)}{(2x+1)(2x-1)}$$

$$58. \frac{2x-1}{4x+1} - \frac{2x}{4x-1} = \frac{(2x-1)(4x-1)}{(4x+1)(4x-1)} - \frac{2x(4x+1)}{(4x+1)(4x-1)} = \frac{(8x^2-6x+1)-(8x^2+2x)}{(4x+1)(4x-1)} = \frac{1-8x}{(4x+1)(4x-1)}$$

$$59. \frac{-x}{x+2} + \frac{x-2}{x} - \frac{x}{x-2} = \frac{-x^2(x-2)}{x(x+2)(x-2)} + \frac{(x-2)^2(x+2)}{x(x+2)(x-2)} - \frac{x^2(x+2)}{x(x+2)(x-2)}$$

$$= \frac{(-x^3+2x^2)}{x(x+2)(x-2)} + \frac{(x^3-2x^2-4x+8)}{x(x+2)(x-2)} + \frac{-(x^3+2x^2)}{x(x+2)(x-2)}$$

$$= \frac{-x^3-2x^2-4x+8}{x(x+2)(x-2)} = -\frac{x^3+2x^2+4x-8}{x(x+2)(x-2)}$$

$$60. \frac{3x}{x-1} + \frac{x+1}{x} - \frac{2x}{x+1} = \frac{3x^2(x+1)}{x(x-1)(x+1)} + \frac{(x+1)^2(x-1)}{x(x-1)(x+1)} - \frac{2x^2(x-1)}{x(x-1)(x+1)}$$

$$= \frac{3x^3+3x^2}{x(x-1)(x+1)} + \frac{x^3+x^2-x-1}{x(x-1)(x+1)} - \frac{2x^3-2x^2}{x(x-1)(x+1)} = \frac{2x^3+6x^2-x-1}{x(x-1)(x+1)}$$

In exercises 61–68, to find the LCD, first factor each denominator completely, form a product of the distinct irreducible factors of each polynomial, and then attach to each factor in this product the greatest exponent that appears on this factor in any of the factored denominators.

$$61. \text{ The denominators are } 3x-6=3(x-2) \text{ and } 4x-8=4(x-2) \Rightarrow \text{LCD} = 3 \cdot 4(x-2) = 12(x-2)$$

$$62. \text{ The denominators are } 7+21x=7(1+3x) \text{ and } 3+9x=3(1+3x) \Rightarrow \text{LCD} = 7 \cdot 3(1+3x) = 21(1+3x)$$

$$63. \text{ The denominators are } 4x^2-1=(2x+1)(2x-1) \text{ and } (2x+1)^2=(2x+1)(2x+1) \Rightarrow \text{LCD} = (2x-1)(2x+1)^2$$

$$64. \text{ The denominators are } (3x-1)^2=(3x-1)(3x-1) \text{ and } 9x^2-1=(3x-1)(3x+1) \Rightarrow \text{LCD} = (3x-1)^2(3x+1)$$

$$65. \text{ The denominators are } x^2+3x+2=(x+1)(x+2) \text{ and } x^2-1=(x-1)(x+1) \Rightarrow \text{LCD} = (x-1)(x+1)(x+2)$$

$$66. \text{ The denominators are } x^2-x-6=(x-3)(x+2) \text{ and } x^2-9=(x-3)(x+3) \Rightarrow \text{LCD} = (x-3)(x+3)(x+2)$$

$$67. \text{ The denominators are } x^2-5x+4=(x-1)(x-4) \text{ and } x^2+x-2=(x-1)(x+2) \Rightarrow \text{LCD} = (x-1)(x-4)(x+2)$$

68. The denominators are  $x^2 - 2x - 3 = (x - 3)(x + 1)$  and  
 $x^2 + 3x + 2 = (x + 1)(x + 2) \Rightarrow \text{LCD} = (x - 3)(x + 1)(x + 2)$

For exercises 69–84, remember to find the LCD first, and then rewrite each rational expression as a rational expression with the LCD as the denominator. Next add or subtract numerators, following the rules for order of operations. Finally simplify the final rational expression. Leave your answer in factored form.

$$69. \frac{5}{x-3} + \frac{2x}{x^2-9} = \frac{5}{x-3} + \frac{2x}{(x-3)(x+3)} = \frac{5(x+3)}{(x-3)(x+3)} + \frac{2x}{(x-3)(x+3)} = \frac{5x+15+2x}{(x-3)(x+3)} = \frac{7x+15}{(x-3)(x+3)}$$

$$70. \frac{3x}{x-1} + \frac{x}{x^2-1} = \frac{3x}{x-1} + \frac{x}{(x-1)(x+1)} = \frac{3x(x+1)}{(x-1)(x+1)} + \frac{x}{(x-1)(x+1)} = \frac{3x^2+3x+x}{(x-1)(x+1)} = \frac{3x^2+4x}{(x-1)(x+1)}$$

$$= \frac{x(3x+4)}{(x-1)(x+1)}$$

$$71. \frac{2x}{x^2-4} - \frac{x}{x+2} = \frac{2x}{(x-2)(x+2)} - \frac{x}{x+2} = \frac{2x}{(x-2)(x+2)} - \frac{x(x-2)}{(x+2)(x-2)}$$

$$= \frac{2x - (x^2 - 2x)}{(x-2)(x+2)} = \frac{-x^2 + 4x}{(x-2)(x+2)} = \frac{x(4-x)}{(x-2)(x+2)}$$

$$72. \frac{3x-1}{x^2-16} - \frac{2x+1}{x-4} = \frac{3x-1}{(x-4)(x+4)} - \frac{2x+1}{x-4} = \frac{3x-1}{(x-4)(x+4)} - \frac{(2x+1)(x+4)}{(x-4)(x+4)}$$

$$= \frac{3x-1 - (2x^2+9x+4)}{(x-4)(x+4)} = \frac{-2x^2-6x-5}{(x-4)(x+4)} = -\frac{2x^2+6x+5}{(x-4)(x+4)}$$

$$73. \frac{x-2}{x^2+3x-10} + \frac{x+3}{x^2+x-6} = \frac{x-2}{(x-2)(x+5)} + \frac{x+3}{(x-2)(x+3)} = \frac{1}{x+5} + \frac{1}{x-2}$$

$$= \frac{x-2}{(x+5)(x-2)} + \frac{x+5}{(x+5)(x-2)} = \frac{2x+3}{(x-2)(x+5)}$$

$$74. \frac{x+3}{x^2-x-2} + \frac{x-1}{x^2+2x+1} = \frac{x+3}{(x-2)(x+1)} + \frac{x-1}{(x+1)(x+1)} = \frac{(x+3)(x+1)}{(x-2)(x+1)(x+1)} + \frac{(x-1)(x-2)}{(x+1)(x+1)(x-2)}$$

$$= \frac{(x^2+4x+3) + (x^2-3x+2)}{(x-2)(x+1)(x+1)} = \frac{2x^2+x+5}{(x-2)(x+1)^2}$$

$$75. \frac{2x-3}{9x^2-1} + \frac{4x-1}{(3x-1)^2} = \frac{2x-3}{(3x-1)(3x+1)} + \frac{4x-1}{(3x-1)^2} = \frac{(2x-3)(3x+1)}{(3x+1)(3x-1)^2} + \frac{(4x-1)(3x+1)}{(3x+1)(3x-1)^2}$$

$$= \frac{(6x^2-11x+3) + (12x^2+x-1)}{(3x+1)(3x-1)^2} = \frac{18x^2-10x+2}{(3x+1)(3x-1)^2} = \frac{2(9x^2-5x+1)}{(3x+1)(3x-1)^2}$$

$$76. \frac{3x+1}{(2x+1)^2} + \frac{x+3}{4x^2-1} = \frac{3x+1}{(2x+1)^2} + \frac{x+3}{(2x-1)(2x+1)} = \frac{(3x+1)(2x-1)}{(2x+1)^2(2x-1)} + \frac{(x+3)(2x+1)}{(2x-1)(2x+1)^2}$$

$$= \frac{(6x^2-x-1) + (2x^2+7x+3)}{(2x-1)(2x+1)^2} = \frac{8x^2+6x+2}{(2x-1)(2x+1)^2} = \frac{2(4x^2+3x+1)}{(2x-1)(2x+1)^2}$$

$$77. \frac{x-3}{x^2-25} - \frac{x-3}{x^2+9x+20} = \frac{x-3}{(x-5)(x+5)} - \frac{x-3}{(x+4)(x+5)} = \frac{(x-3)(x+4)}{(x-5)(x+5)(x+4)} - \frac{(x-3)(x-5)}{(x+4)(x+5)(x-5)}$$

$$= \frac{(x^2+x-12) - (x^2-8x+15)}{(x-5)(x+5)(x+4)} = \frac{9x-27}{(x-5)(x+5)(x+4)} = \frac{9(x-3)}{(x-5)(x+5)(x+4)}$$

$$\begin{aligned}
 78. \quad \frac{2x}{x^2-16} - \frac{2x-7}{x^2-7x+12} &= \frac{2x}{(x-4)(x+4)} - \frac{2x-7}{(x-4)(x-3)} = \frac{2x(x-3)}{(x-4)(x+4)(x-3)} - \frac{(2x-7)(x+4)}{(x-4)(x+4)(x-3)} \\
 &= \frac{(2x^2-6x)-(2x^2+x-28)}{(x-4)(x+4)(x-3)} = \frac{-7x+28}{(x-4)(x+4)(x-3)} \\
 &= \frac{-7(x-4)}{(x-4)(x+4)(x-3)} = -\frac{7}{(x+4)(x-3)}
 \end{aligned}$$

$$\begin{aligned}
 79. \quad \frac{3}{x^2-4} + \frac{1}{2-x} - \frac{1}{2+x} &= -\frac{3}{4-x^2} + \frac{1}{2-x} - \frac{1}{2+x} = -\frac{3}{(2-x)(2+x)} + \frac{1}{2-x} - \frac{1}{2+x} \\
 &= \frac{-3}{(2-x)(2+x)} + \frac{1(2+x)}{(2-x)(2+x)} - \frac{1(2-x)}{(2+x)(2-x)} = \frac{-3+(2+x)-(2-x)}{(2-x)(2+x)} = \frac{2x-3}{(2-x)(2+x)}
 \end{aligned}$$

$$\begin{aligned}
 80. \quad \frac{2}{5+x} + \frac{5}{x^2-25} + \frac{7}{5-x} &= \frac{2}{5+x} - \frac{5}{25-x^2} + \frac{7}{5-x} = \frac{2}{5+x} - \frac{5}{(5-x)(5+x)} + \frac{7}{5-x} \\
 &= \frac{2(5-x)}{(5+x)(5-x)} - \frac{5}{(5-x)(5+x)} + \frac{7(5+x)}{(5-x)(5+x)} \\
 &= \frac{(10-2x)-5+(35+7x)}{(5+x)(5-x)} = \frac{5x+40}{(5+x)(5-x)} = \frac{5(x+8)}{(5+x)(5-x)}
 \end{aligned}$$

$$81. \quad \frac{x+3a}{x-5a} - \frac{x+5a}{x-3a} = \frac{(x+3a)(x-3a)}{(x-5a)(x-3a)} - \frac{(x+5a)(x-5a)}{(x-3a)(x-5a)} = \frac{(x^2-9a^2)-(x^2-25a^2)}{(x-5a)(x-3a)} = \frac{16a^2}{(x-5a)(x-3a)}$$

$$82. \quad \frac{3x-a}{2x-a} - \frac{2x+a}{3x+a} = \frac{(3x-a)(3x+a)}{(2x-a)(3x+a)} - \frac{(2x+a)(2x-a)}{(3x+a)(2x-a)} = \frac{(9x^2-a^2)-(4x^2-a^2)}{(2x-a)(3x+a)} = \frac{5x^2}{(2x-a)(3x+a)}$$

$$\begin{aligned}
 83. \quad \frac{1}{x+h} - \frac{1}{x} &= \frac{x}{x(x+h)} - \frac{1(x+h)}{x(x+h)} \\
 &= \frac{x-(x+h)}{x(x+h)} = -\frac{h}{x(x+h)}
 \end{aligned}$$

$$\begin{aligned}
 84. \quad \frac{1}{(x+h)^2} - \frac{1}{x^2} &= \frac{x^2}{x^2(x+h)^2} - \frac{1(x+h)^2}{x^2(x+h)^2} \\
 &= \frac{x^2-(x+h)^2}{x^2(x+h)^2} \\
 &= \frac{x^2-(x^2+2xh+h^2)}{x^2(x+h)^2} \\
 &= \frac{-2xh-h^2}{x^2(x+h)^2} = -\frac{h(2x+h)}{x^2(x+h)^2}
 \end{aligned}$$

$$85. \quad \frac{\frac{2}{x}}{\frac{3}{x^2}} = \frac{2}{x} \cdot \frac{x^2}{3} = \frac{2x}{3}$$

$$86. \quad \frac{\frac{-6}{x}}{\frac{2}{x^3}} = \frac{-6}{x} \cdot \frac{x^3}{2} = -3x^2$$

87. Using Method 1:

$$\begin{aligned}
 \frac{\frac{1}{x}}{1-\frac{1}{x}} &= \frac{\frac{1}{x}}{\frac{x-1}{x}} = \frac{\frac{1}{x}}{\frac{x-1}{x}} = \frac{1}{x} \cdot \frac{x}{x-1} \\
 &= \frac{1}{x-1}, \quad x \neq 0, 1
 \end{aligned}$$

88. Using Method 1:

$$\begin{aligned}
 \frac{\frac{1}{x^2}}{\frac{1}{x^2}-1} &= \frac{\frac{1}{x^2}}{\frac{1-x^2}{x^2}} = \frac{\frac{1}{x^2}}{\frac{1-x^2}{x^2}} = \frac{1}{x^2} \cdot \frac{x^2}{1-x^2} \\
 &= \frac{1}{1-x^2} = \frac{1}{(1-x)(1+x)}, \quad x \neq -1, 0, 1
 \end{aligned}$$

89. Using Method 2:

$$\frac{\frac{1}{x}-1}{\frac{1}{x}+1} = \frac{\left(\frac{1}{x}-1\right)x}{\left(\frac{1}{x}+1\right)x} = \frac{1-x}{1+x}, \quad x \neq 0, x \neq -1$$